

Anticipated Classification
of this Application:
Class _____ Subclass _____

06-02-00
Prior Application:
Examiner S. McGarry
Group Art Unit 1635

Attorney's
Docket
No. 50995-B/JPW/EMW

June 1, 2000

HONORABLE ASSISTANT COMMISSIONER FOR PATENTS
Washington, D.C. 20231

06/01/00
JC822 U.S. PTO
R:

This is a request for filing a X CONTINUATION

DIVISIONAL _____ CONTINUATION-IN-PART application under

X 37 C.F.R. §1.53(b) _____ 37 C.F.R. §1.53(d), of pending prior application

Serial No. 08/654,482 filed on May 28, 1996 of

Riccardo Dalla-Favera

for

Inventor(s)

IDENTIFICATION OF GENES ALTERED IN MULTIPLE MYELOMA

Title of Invention

- 06/01/00
JC822 U.S. PTO
R:
1. X Enclosed is a copy of the prior application, as originally filed and an affidavit or declaration verifying it as a true copy.
 2. X A verified statement to establish small entity status under 37 C.F.R. §1.9 and 1.27
_____ is enclosed.
X was filed in the prior application and such status is still proper and desired (37 C.F.R. §1.28(a)).
 3. X The filing fee is calculated as follows:

CLAIMS AS FILED, LESS ANY CLAIMS CANCELLED BY AMENDMENT

	NUMBER FILED		NUMBER EXTRA*		RATE		FEE		
					SMALL ENTITY	OTHER ENTITY		SMALL ENTITY	OTHER ENTITY
Total Claims	7 -20	=	0	X	\$ 9.00	\$18.00	=	\$ 0	\$
Independent Claims	2 -3	=	0	X	\$39.00	\$78.00	=	\$ 0	\$
Multiple Dependent Claims Presented: ___ Yes <u>X</u> No					\$130.00	\$260.00	=	\$ 0	\$
*If the different in Col. 1 is less than zero, enter "0" in Col. 2					BASIC FEE			\$ 345	\$ 690
					TOTAL FEE			\$345.00	\$

4. X The Commissioner is hereby authorized to charge payment of the following fees associated with this application or credit any overpayment to Deposit Account No. 03-3125.

 X Any additional filing fees required under 37 C.F.R. §1.16.

 X Any patent application processing fees under 37 C.F.R. §1.17.

 The issue fees set forth in 37 C.F.R. §1.18 at or before mailing of the Notice of Allowance, pursuant to 37 C.F.R. §1.311(b).

5. X Three copies of this sheet are enclosed.

6. X A check in the amount of \$ 345.00 is enclosed.

7. Cancel claims .

8. Amend the specification by inserting before the first line the sentence: --This is a continuation division of application Serial No. , filed .--

9. X 21 Sheet(s) of informal X formal drawing(s) is/are enclosed.

10. Transfer the drawings from the prior application to this application and abandon said prior application as of the filing date accorded this application. A duplicate copy of this sheet is enclosed for filing in the prior application file.

11. Priority of application No. filed on in (country) is claimed under 37 U.S.C. §119.

 The certified copy of the priority application has been filed in prior application Serial No. , filed .

12. X The prior application is assigned of record to The Trustees of Columbia University in the City of New York. (a copy of the Assignment is attached)

13. X A preliminary amendment is enclosed.

14. X The power of attorney in the prior application is to:

John P. White (Reg. No. 28,678); Thomas F. Moran (Reg. No. 16,579); Norman H. Zivin (Reg. No. 25,385); Ivan S. Kavrukov (Reg. No. 25,161); Christopher C. Dunham (Reg. No. 22,031); Thomas G. Carulli (Reg. No. 30,616); Robert D. Katz (Reg. No. 30,141); Peter J. Phillips (Reg. No. 29,691); Richard S. Milner (Reg. No. 33,970); Albert Wai-Kit Chan (Reg. No. 36,479); Kristina L. Konstas (Reg. No. 37,864); Mary Anne P. Tanner (Reg. No. 40,197); Timothy X. Witkowski (Reg. No. 40,232); and Mary Catherine DiNunzio (Reg. No. 37,306)

Applicants: Riccardo Dalla-Favera
U.S. Serial No.: Not Yet Known
Filed: Herewith
Cont/Div.
Page 3

June 1, 2000

- (a) X The power appears in the original papers in the prior application.
(a copy of the Declaration is attached)
- (b) Since the power does not appear in the original papers, a copy of the power in the prior application is enclosed.
- (c) X Address all future communications to:
(May only be completed by applicant,
or attorney or agent of record.)

John P. White

Cooper & Dunham LLP


1185 Avenue of the Americas

New York, New York 10036

15. X Also enclosed Express Mail Certificate of Mailing No. EJ 807 507 669 US, a loose set of loose figures, Statement in Accordance and a computer diskette of Sequence Listing.
16. X I hereby verify that the attached papers are a true copy of prior application Serial No. 08/654,482 as originally filed on May 28, 1996.

The undersigned declares further that all statements made herein of his own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statement and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

June 1, 2000
Date


Signature John P. White, Reg. No. 28,678

 INVENTOR(S)
 ASSIGNEE OF COMPLETE INTEREST
 X ATTORNEY OR AGENT OF RECORD
 FILED UNDER 37 C.F.R. §1.34(a)

Address of Signator:

Cooper & Dunham LLP

1185 Avenue of the Americas

New York, New York 10036

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Riccardo Dalla-Favera
Serial No.: Not Yet Known (Continuation of U.S. Serial No.
08/654,482, filed May 28, 1996)
Filed: Herewith
For: IDENTIFICATION OF GENES ALTERED IN MULTIPLE MYELOMA

1185 Avenue of the Americas
New York, New York 10036
June 1, 2000

Assistant Commissioner for Patents
Washington, D.C. 20231
Box: Patent Application

SIR:

PRELIMINARY AMENDMENT

Please amend the above-identified application as follows:

In the specification:

At page 1, after the title and before line 3, please insert:

--This application is a continuation of U.S. Serial No. 08/654,482,
filed May 26, 1996, now allowed, the contents of which are hereby
incorporated by reference.--

In the claims:

Please cancel claims 1-87, 93-97 and 99-101 without prejudice to
applicant's right to pursue the subject matter of these claims in

Applicant: Riccardo Dalla-Favera
Serial No.: Not Yet Known(Continuation of U.S. Serial No.
08/654,482, filed May 28, 1996)
Filed: Herewith

a future continuation application.

Please amend claims 89-92 and 98 under 37 C.F.R. §1.121(b) by deleting the bracketed materials and inserting the underlined materials as follows:

--89.(Amended) A purified human MUM-1 protein of claim 88[.],wherein the MUM-1 protein has the same amino acid sequence as shown in Figure 5B (SEQ. ID NO:14).--

--90.(Amended) An antibody directed to [a] the purified MUM-1 protein of claim 89.--

--91.(Amended) An antibody capable of specifically recognizing MUM-1 protein, wherein the MUM-1 protein has the same amino acid sequence as shown in Figure 5B (SEQ. ID NO:14).--

--92.(Amended) [An] The antibody of claim 91, wherein the MUM-1 protein is a human MUM-1 protein.--

--98.(Amended) [An] A monoclonal antibody of [any one of claims] claim 90._, 91 and 92.]--

Please add new claims 102-103 as follows:

Applicant: Riccardo Dalla-Favera
Serial No.: 08/654,482
Filed: May 28, 1996
Page 3

--102. (New) A monoclonal antibody of claim 91.--

--103. (New) A monoclonal antibody of claim 92.--

REMARKS

Claims 1-101 were pending in the subject application. Applicant has hereinabove canceled claims 1-87, 93-97 and 99-101 without disclaimer or prejudice to applicant's right to pursue the subject matter of these claims at a later date. Applicant has also amended claims 89-92 and 98 and added new claims 102-103. Support for amended claims 89 and 91 may be found in the specification inter alia on page 21, lines 2-4 and Figure 5B. Support for amended claim 92 may be found in the specification inter alia on page 32, lines 10-12 and page 21, lines 2-4. Support for amended claim 98 may be found in the specification inter alia on page 32, lines 22-23. Support for new claims 102-103 may be found in the specification inter alia on page 32, lines 23-27 and in previously pending claim 98. Applicant maintains that the amendments are fully supported by the specification and do not raise any issue of new matter. Accordingly, applicant respectfully requests that the Examiner enter the Amendment. Upon entry of the Amendment, claims 88-92 and 98, as amended, and new claims 102-103 will be pending and under examination.


If a telephone interview would be of assistance in advancing prosecution of the subject application, applicant's undersigned attorney invites the Examiner to telephone at the number provided

Applicant: Riccardo Dalla-Favera
Serial No.: 08/654,482
Filed: May 28, 1996
Page 4

below.

No fee is deemed necessary in connection with the filing of this Preliminary Amendment. However, if any additional fee is required, authorization is hereby given to charge the amount of such fee to Deposit Account No. 03-3125.

Respectfully submitted



John P. White
Registration No. 28,678
Attorney for the Applicant
Cooper & Dunham LLP
1185 Avenue of the Americas
New York, New York 10036
(212) 278-0400

**Application
for
United States Letters Patent**

To all whom it may concern:

Be it known that I, Riccardo Dalla-Favera

have invented certain new and useful improvements in

IDENTIFICATION OF GENES ALTERED IN MULTIPLE MYELOMA

of which the following is a full, clear and exact description.

IDENTIFICATION OF GENES ALTERED IN MULTIPLE MYELOMA

The invention disclosed herein was made with Government support under NIH Grant No. CA 44025. Accordingly, the U.S. Government has certain rights in this invention.

Background of the Invention

Throughout this application, various references are referred to within parentheses. Disclosures of these publications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which this invention pertains. Full bibliographic citation for these references may be found at the end of this application, preceding the claims.

Multiple myeloma (MM) is an incurable B cell tumor affecting B cell end-stage differentiation. Clinically, the course of MM is similar to end-stage plasma cell leukemia (PCL), i.e., there is an uncontrollable proliferation of myeloma cells accompanied by numerous complications, including hyperviscosity syndromes, hypercalcemia, infections, multiple bone fractures, and organ failure.

Non-random chromosomal translocation is known to play a crucial role in the tumorigenesis of hematologic malignancies (1). In B-cell lymphomas, many important proto-oncogenes deregulated by juxtaposition to immunoglobulin (Ig) gene locus have been identified. Each proto-oncogene is associated with a specific subtype of lymphoma, such as c-MYC in Burkitt's lymphoma, Cyclin D1 in mantle cell lymphoma, BCL-2 in follicular lymphoma and BCL-6 in diffuse large cell lymphoma (2-8). In contrast, little is known about molecular alterations of

human MM/PCL, due to the difficulty in cytogenetic analysis. However, previous cytogenetic reports have shown a 14q+ chromosome, suggesting the existence of a chromosomal translocation involving the Ig heavy chain (IgH) locus, is observed in 20 ~ 30 % of the MM/PCL cases and it is the most frequent consistent abnormality (9-12). Even in such cases, most cytogenetic data have failed to identify donor chromosomes other than 11q13, 8q24, and 18q21, where proto-oncogenes Cyclin D1, BCL-2, and c-MYC are located, respectively. Among them, the 11q13 locus has been demonstrated to be involved in nearly 5~10% of the cases and also in 62% of the established cell lines (13). The t(11;14)(q13;q32) translocation is also accompanied by a corresponding overexpression of the Cyclin D1 gene, which raises a strong possibility of the involvement of this gene, although the breakpoints at 11q13 do not cluster like those of the lymphoma cases (14-16). Recent advances in fluorescence in situ hybridization (FISH) have made it possible to clarify both the frequency of the 14q+ chromosomes and the partner chromosomes of the IgH loci. One such report revealed an intriguing result, i.e., that numerous chromosomal loci are able to translocate to IgH locus, including 6p21, 1q21, 3p11, 7q11, 11q23 (17). This has prompted a search for the proto-oncogenes deregulated by the regulatory elements of the IgH gene for a further understanding of the molecular mechanisms of MM/PCL. In the present study, one candidate proto-oncogene, MUM1 (multiple myeloma oncogene 1), was found juxtaposed to the IgH gene as a result of t(6;14)(p25; q32) translocation in human myeloma cell line, SKMM-1. Over expression of the MUM1 mRNA was observed in this cell line. A second gene, called MUM-2 was found translocated in proximity to the IgH gene on

chromosome 14q32 in human myeloma cell line, U-266.

The method of analysis of 14q+ chromosomal translocations and identification of the genes altered in multiple myeloma of this invention are useful since 1) no method is currently available to determine the chromosomal sequences involved in 14q+ translocations, the most important cytogenetic lesions associated with MM pathogenesis; 2) no specific gene lesion is currently known for MM; 3) no diagnostic method based on gene/DNA lesion is currently available for MM and 4) there are no therapeutic approaches aimed at counteracting the action of abnormal gene products in MM.

Summary of the Invention

This invention provides a method of determining a chromosomal breakpoint in a subject suffering from multiple myeloma which comprises steps of: (a) obtaining a DNA sample from the subject suffering from multiple myeloma; (b) determining whether there is J and C disjunction in the immunoglobulin heavy chain gene in the obtained DNA sample; (c) obtaining a genomic library having clones which contain genomic DNA fragments from the DNA sample which shows positive J and C disjunction; (d) selecting and isolating clones of the obtained library which show positive hybridization with a probe which is capable of specifically hybridizing with the C but not the J region of the immunoglobulin heavy chain gene; (e) preparing fluorescent probes from the genomic DNA fragments of the isolated clones from step (d); (f) hybridizing said fluorescent probes with metaphase chromosomes; and (g) determining the identity of the chromosomes which are capable of hybridizing to said fluorescent probes, wherein the identification of a chromosome other than chromosome 14 would indicate that the chromosomal breakpoint is between chromosome 14 and the identified chromosome, thereby determining a chromosomal breakpoint in a subject suffering from multiple myeloma.

This invention provides a method to identify a gene other than the immunoglobulin gene which is located in chromosome 14, altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma which comprises steps of: a) selecting a probe having a sequence of a chromosome other than chromosome 14, identified at the chromosomal breakpoint detected in a subject suffering from multiple myeloma, wherein said probe is capable of hybridizing to the unique sequence of the gene other than

the immunoglobulin gene altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma; b) contacting said probe with mRNA isolated from a cell under conditions permitting formation of a complex between said probe and the mRNA; c) isolating the complex resulting from step (b); d) determining the sequence of the mRNA in the isolated complex, thereby determining the identity of the gene.

10 This invention provides a gene designated *MUM-1*. This invention provides a gene designated *MUM-2*. This invention provides an isolated nucleic acid molecule encoding a MUM protein. This invention provides a DNA encoding a MUM protein. This invention provides a cDNA encoding a MUM protein. This invention provides a genomic DNA molecule encoding a MUM protein. This invention provides a RNA molecule encoding a MUM protein. This invention provides an isolated nucleic acid molecule encoding a human *MUM-1* protein. This invention provides an isolated nucleic acid molecule encoding a human *MUM-2* protein. This invention provides an isolated nucleic acid molecule encoding a MUM protein operatively linked to a promoter of RNA transcription. This invention provides a vector comprising the an isolated cDNA encoding a MUM protein. This invention provides a vector which comprises an isolated cDNA encoding a MUM protein. This invention provides a vector which comprises an isolated cDNA encoding a MUM protein, wherein the vector is a plasmid. This invention provides a host cell for the vector which comprises an isolated cDNA encoding a MUM protein.

This invention provides a nucleic acid probe comprising a

nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a MUM protein. This invention provides a nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding a MUM protein.

This invention provides a nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding a MUM protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

This invention provides a nucleic acid probe comprising a the sequence of a nucleic acid molecule encoding a MUM-1 protein which is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14.

This invention provides a nucleic acid probe comprising a the sequence of a nucleic acid molecule encoding a MUM-2 protein which is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14.

This invention provides a method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-1 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-1 protein. This invention provides a method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-2 protein in a sample from a subject which comprises detecting in a sample from the

subject a rearrangement of nucleic acid encoding MUM-2 protein.

5 This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to an mRNA molecule encoding a human MUM-1 protein so as to prevent overexpression of the mRNA molecule. This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to an mRNA molecule
10 encoding a human MUM-2 protein so as to prevent overexpression of the mRNA molecule.

15 This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to an isolated cDNA molecule encoding a MUM protein. This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to the isolated genomic DNA molecule encoding a MUM protein. This invention provides an antisense oligonucleotide having a sequence
20 capable of specifically hybridizing to an isolated RNA molecule encoding a MUM protein.

25 This invention provides a purified MUM protein. This invention provides a purified MUM-1 protein. This invention provides an antibody directed to a purified MUM-1 protein. This invention provides an antibody capable of specifically recognizing MUM-1 protein. This invention provides a purified MUM-2 protein. This invention provides an antibody directed to a purified MUM-2 protein. This invention
30 provides an antibody capable of specifically recognizing a MUM-2 protein.

This invention provides a pharmaceutical composition

comprising an amount of an oligonucleotide effective to prevent overexpression of a human MUM-1 protein and a pharmaceutically acceptable carrier capable of passing through a cell membrane. This invention provides a
5 pharmaceutical composition comprising an amount of the oligonucleotide effective to prevent overexpression of a human MUM-2 protein and a pharmaceutically acceptable carrier capable of passing through a cell membrane.

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219

Brief Description of the Figures

Figure 1. JH-C μ dissociation in *Bam*HI digested DNA of the 14q+ SK-MM-1 cell line. A 10 μ g of the high molecular weight DNA was completely digested with *Bam*HI, loaded on each lane and blotted. The same filter was sequentially hybridized with JH, C μ , Cy2, and 0.7B/H probes. JH probe detects two rearranged bands of 12.0 kb and 9.7kb. The 9.7 kb band is comigrated with that probed with Cy2 probe, suggesting it to be a physiological rearrangement. On the other hand, one allele of the C μ locus is deleted and another is rearranged (6.5 kb) without being comigrated with rearranged bands of JH. Therefore, 12.0 kb and 6.5 kb bands detected by JH and C μ (shown by arrowheads) might represent unknown derivative chromosome and derivative 14 chromosome, respectively. As expected, 0.7B/H probe (Fig. 2A) detected the rearranged band comigrated with 6.5kb band of C μ . Dashed lines show the comigration. Size markers of λ /*Hind*III are shown on the left.

Figures 2A-B. Molecular cloning of the breakpoints of the t(6;14) translocation and germline walking at MUM1 locus. (A) Restriction maps of λ SKB-4a and λ SKS-3 clones representing derivative 6 and 14 are shown, together with germline maps of IgH locus at 14q32 and MUM1

locus at 6p25. Arrows indicate the chromosomal breakpoints. B, *Bam*HI; E, *Eco*RI; H, *Hind*III. (B) Comparison of the nucleotide sequences around the breakpoints on derivative 6 and derivative 14 chromosome. Homologous regions are indicated by dashes. The arrow indicates the breakpoint. Nucleotide numbers shown below are the same as in the Sp sequence reported by Sun, et al. (18).

Figure 3. Mapping of the MUM1 locus to chromosome 6p25. λ MUM-3 genomic clone (Figure 2A) was used as a probe for *in situ* hybridization. The white arrow indicates the fluorescence signal on chromosome 6 band p25. Right panel shows the G-banding picture stained with DAPI.

Figures 4A-C. Expression of the MUM1 gene in hematopoietic lineage. A 10 ug aliquot of total RNA was loaded on each lane and Northern blot analysis was performed using the 2.1H probe (Figure 2A). GAPDH or β -actin probes were used to control for amount of RNA loaded. (A) MUM1 RNA expression in various hematopoietic cell lines. MUM1 RNA is detected in B cell and mature T cell lines as a single 6kb transcript. HELA, epithelial lineage; LCL, Epstein-Barr virus-transformed lymphoblastoid cell line; RAMOS and SK-MM-1, B-cell lineage; HUT-78 and MOLT-4, T-cell

lineage; HL-60 and U937, myelomonocytic lineage; K562, erythroid lineage. Dashes indicate 28S and 18S. (B) Expression in B cell lines derived from various stages of B cell differentiation. MUM1 RNA is seen throughout the B cell development except for BJAB cell line. 697, pre-B cell stage; RAMOS and BJA-B, Burkitt cell line representing mature-B cell stage; RPMI-8226 and U-266, plasma cell stage. (C) Comparison of the expression level among myeloma cell lines. MUM1 RNA is overexpressed in SK-MM-1 cell line carrying t(6;14). Overexpression of the MUM1 is also demonstrated in XG-4, XG-7, and XG-10 cell lines. RPMI-8226, U-266, EJM, and SKMM-1 are IL-6 (interleukin-6) independent lines, whereas XG-1, XG-2, XG-4, XG-5, XG-6, XG-7, and XG-10 are IL-6 dependent lines.

Figures 5A-B. Sequence of MUM1 cDNA and structure of its predicted protein product. (A) Restriction map of the MUM1 cDNA and the position of the open reading frame (box). The solid box indicates approximate position of the DNA binding domain. Sc, *SacII*; A, *ApaI*; P, *PstI*; H, *HindIII*; S, *SacI* (B) Nucleotide sequence of the MUM1 cDNA and corresponding amino acid sequence. Putative translation initiation codons and preceding stop codons appearing in frame are underlined. The asterisk indicates the translation stop

codon.

Figures 6A-B. Homology between MUM1 and other IRF family proteins. (A) Similarity at N-terminal DNA binding domain. Black background indicates identical residues found more than four times. Gray indicates conserved residues that appear in at least four sequences at a given position. Conserved tryptophan residues in DNA binding domain among IRF family members are indicated by closed circles. (B) Similarity at C-terminal region between human MUM1, Mouse LSIRF/Pip, Human ICSBP, Human ISGF3 γ , and Human IRF-3. Black and gray background are as in (A).

Figure 7. Genomic organization of the MUM1 gene and location of the chromosomal breakpoints in multiple myeloma. Filled boxes indicate the coding regions and empty boxes indicate the noncoding regions. The position, and the size of each exon of the MUM1 gene are approximate and have been determined by the hybridizations. One exon in each restriction fragment may consist of more than two exons. Translation initiation codon (ATG) and stop codon (TGA) are indicated. Genomic probes used for further investigations are shown as solid bars below the map. Arrows indicate the chromosomal breakpoints of SKMM-1 cell line and case 10. B, *Bam*HI; E, *Eco*RI; H, *Hind*III.

Figure 8. Scheme of the t(6;14)(p25;q32) translocation involving the MUM1 and the immunoglobulin heavy chain (IgH) gene loci. VH-D-J-CH indicates variable-diversity joining-constant region of the IgH gene. Direction of the MUM1 gene on the chromosome 6 is tentatively drawn.

Figure 9A-B. Demonstration of JH-Cα disjunction in U-266 cells and cloning of normal and 14q+ chromosomal breakpoints. (A) The panel shows the results of Southern blot analysis of BamHI digested U-266 and normal control (placenta) genomic DNA using the indicated JH and Cα probes. The arrowheads indicate two DNA fragments containing Cα sequences not linked to JH sequences, suggesting the presence of a chromosomal breakpoint in 14q32. (B) The panel provides a schematic representation of the phage clones isolated from a library constructed from U-266 DNA and screened with a Cα probe. Based on restriction enzyme analysis, the three cloned regions represent a normal Cα region (14q32 germ-line), and two rearranged regions (der.14 and 14q32) containing unknown sequences linked to Cα sequences. The 2.5BE probe used for Northern blot analysis of MUM2 transcripts (Fig. 10) is also shown.

Figure 10. Identification of MUM2 RNA transcripts. The

figure shows the results of a Northern blot analysis of RNA extracted from various MM/PCL cell lines using the 2.5BE probe (see Fig. 9) or GAPDH probe (as a control for RNA loading). A 1.9 Kb RNA transcript is detectable in some cell lines including U-266, indicating that the 2.5BE fragments represents part of a gene, MUM2.

Figure 11A-B. Schematic representation of IgH DNA rearrangements in normal B cells and in tumors carrying chromosomal translocations breaking the S region of the IgH locus. Note that in physiological IgH rearrangements (panel 11A) JH sequences and C sequences (C_μ before and C_γ after switch recombination, respectively) are consistently found within the same BamHI restriction fragment. Conversely, JH and C sequences are not linked, and are present on two different chromosomes [derivative X and derivative 14(14q+)] in cells carrying a chromosomal translocation breaking the switch region (panel 11B)

Figure 12A-B. MUM1 cDNA. cDNA insert is cloned into EcoRI/BamHI site of the pBluescript KS+. Bacteria strain used is DH5α cells. pcMUM1.16a contains full length open reading frame of nt. 217-1572.

Figure 13. Breakpoint Cloning of the U-266 Cell Line.

-15-

pMUM2-8 has a 22.0 KB insert in BamHI site
of pBluescript KS+.

Detailed Description of the Invention

The following standard abbreviations are used throughout the specification to indicate specific nucleotides:

5

C=cytosine A=adenosine
T=thymidine G=guanosine

10

This invention provides a method of determining a chromosomal breakpoint in a subject suffering from multiple myeloma which comprises steps of: (a) obtaining a DNA sample from the subject suffering from multiple myeloma; (b) determining whether there is J and C disjunction in the immunoglobulin heavy chain gene in the obtained DNA sample;

15

(c) obtaining a genomic library having clones which contain genomic DNA fragments from the DNA sample which shows positive J and C disjunction; (d) selecting and isolating clones of the obtained library which show positive hybridization with a probe which is capable of specifically hybridizing with the C but not the J region of the immunoglobulin heavy chain gene; (e) preparing fluorescent probes from the genomic DNA fragments of the isolated clones

20

from step (d); (f) hybridizing said fluorescent probes with metaphase chromosomes; and (g) determining the identity of the chromosomes which are capable of hybridizing to said fluorescent probes, wherein the identification of a chromosome other than chromosome 14 would indicate that the chromosomal breakpoint is between chromosome 14 and the identified chromosome, thereby determining a chromosomal breakpoint in a subject suffering from multiple myeloma.

25

30

In an embodiment, step (b) of the above described method of this invention is performed by Southern blotting. In another embodiment, step (b) of the above method of this

invention is performed by polymerase chain reaction (PCR) with appropriate probes. Polymerase chain reaction is well known in the art. Since the sequences of both the C and J regions of an immunoglobulin heavy chain gene are known,
5 appropriate probes for PCR may routinely be designed.

In an embodiment, the genomic library is a phage vector library. In another embodiment, the genomic DNA fragments are generated by cleaving genomic DNA from cells of the
10 subject with an appropriate restriction enzyme. In a further embodiment, the restriction enzyme is *Bam*HI. In an embodiment, the restriction enzyme is *Sau*3AI. In another embodiment, the probe of step (d) is a human IgH J region JH probe. In a further embodiment, the probe of step (d) is a
15 human IgH C μ probe. In an embodiment, the probe of step (d) is a human IgH C γ 2 probe. In another embodiment, the chromosomal breakpoint identified is a t(6;14)(p25;q32) translocation. In an embodiment, the chromosomal breakpoint identified is a t(14;15) translocation.

This invention provides a method to identify a gene other than the immunoglobulin gene which is located in chromosome 14, altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma which comprises
20 steps of: a) selecting a probe having a sequence of a chromosome other than chromosome 14, identified at the chromosomal breakpoint detected in a subject suffering from multiple myeloma, wherein said probe is capable of hybridizing to the unique sequence of the gene other than
25 the immunoglobulin gene altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma; b) contacting said probe with mRNA isolated from a cell under conditions permitting formation of a complex between said probe and the mRNA; c) isolating the complex resulting from
30 step (b); and d) determining the sequence of the mRNA in the
35

isolated complex, thereby determining the identity of the gene.

5 In an embodiment, step (d) of the method to identify a gene other than the immunoglobulin gene which is located in chromosome 14, altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma comprises steps of: i) synthesizing complementary DNA to the mRNA; and ii) performing sequence analysis of the complementary DNA to
10 determine the sequence of the mRNA.

15 This invention provides a gene identified by the method to identify a gene other than the immunoglobulin gene which is located in chromosome 14, altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma.

20 As used herein, "MUM" means any gene rearranged in 14q+ chromosomal abnormalities associated with multiple myeloma.

25 This invention provides a gene identified by the above method designated *MUM-1*. This invention provides a gene identified by the above method designated *MUM-2*.

30 This invention provides a gene identified by the above method, wherein the gene identified comprises a nucleic acid encoding a MUM protein. In an embodiment, the gene identified by the above method comprises a nucleic acid encoding a MUM-1 protein. In another embodiment, the gene identified by the above method comprises a nucleic acid encoding a MUM-2 protein.

This invention provides an isolated nucleic acid molecule encoding a MUM protein. In an embodiment, the isolated

nucleic acid molecule encoding a MUM protein is a DNA molecule. In another embodiment, the isolated nucleic acid molecule encoding a MUM protein is a cDNA molecule.

5

In an embodiment, a cDNA nucleic acid molecule encoding a MUM-1 protein is cloned into a pBluescript KS+ and the resulting plasmid is designated as pcMUM1-1.6a (ATCC Accession No. _____). Plasmid pcMUM1-1.6a was deposited on
10 May 28, 1996 with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. Plasmid pcMUM1-1.6a
15 was accorded ATCC Accession Number _____.

In another embodiment, a partial cDNA nucleic acid molecule encoding a MUM-1 protein is cloned into a pBluescript KS+ and the resulting plasmid is designated as pMUM1-2.4B/N
20 (ATCC Accession No. _____). Plasmid pMUM1-2.4B/N was deposited on May 28, 1996 with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of
25 Microorganisms for the Purposes of Patent Procedure. Plasmid pMUM1-2.4B/N was accorded ATCC Accession Number _____.

In another embodiment, a partial cDNA nucleic acid molecule
30 encoding a MUM-1 protein is cloned into a pBluescript KS+ and the resulting plasmid is designated as pMUM1-7.7B (ATCC Accession No. _____). Plasmid pMUM1-7.7B was deposited on May 28, 1996 with the American Type Culture Collection

(ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. Plasmid pMUM1-7.7B
5 was accorded ATCC Accession Number _____.

In another embodiment, a partial cDNA of the nucleic acid molecule encoding a MUM-2 protein is cloned into a pBluescript KS+ and the resulting plasmid is designated as
10 pMUM2-8 (ATCC Accession No. _____). Plasmid pMUM2-8 was deposited on May 28, 1996 with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of
15 Microorganisms for the Purposes of Patent Procedure. Plasmid pMUM2-8 was accorded ATCC Accession Number _____.

In an embodiment, the isolated DNA molecule encoding a MUM protein is a cDNA molecule having the nucleotide sequence
20 shown in Figure 5B (SEQ. ID NO _____).

In an embodiment, the isolated DNA molecule encoding a MUM protein is genomic DNA molecule. In an embodiment, the isolated nucleic acid molecule encoding a MUM protein is an
25 RNA molecule.

In an embodiment, the isolated nucleic acid encodes a human MUM-1 protein. In another embodiment, the isolated nucleic acid molecule encodes a human MUM-2 protein.

30 In an embodiment, isolated nucleic molecule encodes the a human MUM-1 protein having substantially the same amino acid

sequence as shown in Figure 5B (SEQ. ID NO). In an embodiment, isolated nucleic molecule encodes a human MUM-1 protein having the same amino acid sequence as shown in Figure 5B (SEQ. ID NO). In an embodiment, the isolated
5 nucleic acid molecule encoding a MUM protein is operatively linked to a promoter of RNA transcription.

This invention provides a vector comprising a cDNA molecule encoding a MUM protein. In an embodiment, a vector
10 comprising cDNA encoding for MUM-1 is designated pcMUM1.6a. In an embodiment, a vector comprising partial cDNA encoding for MUM-1 is designated pMUM1.2.4B/N. In an embodiment, a vector comprising partial cDNA encoding for MUM-1 is designated pMUM1-7.7B. In an embodiment, a vector
15 comprising partial cDNA encoding for MUM-2 is designated pMUM2-8. In an embodiment, a vector comprises genomic DNA encoding for MUM. In an embodiment, the vector is a plasmid. In an embodiment, a host cell comprises the vector comprising cDNA encoding for MUM. In an embodiment, a host
20 cell comprises the vector comprising genomic DNA encoding for MUM. In a further embodiment, the host cell comprising vectors comprising cDNA encoding for MUM or comprising genomic DNA encoding for MUM is selected from a group consisting of a bacterial cell, a plant cell, and insect
25 cell and a mammalian cell.

This invention provides a nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a unique sequence included
30 within the sequence of a nucleic acid molecule encoding a MUM protein. This invention provides a nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the

isolated nucleic acid molecule encoding a MUM protein.

As used herein, the phrase "specifically hybridizing" means the ability of a nucleic acid molecule to recognize a nucleic acid sequence complementary to its own and to form double-helical segments through hydrogen bonding between complementary base pairs.

In an embodiment, the nucleic acid probe specifically hybridizes with nucleic acid encoding MUM-1. In an embodiment, the nucleic acid probe is complementary to nucleic acid encoding MUM-1. In an embodiment, the nucleic acid probe specifically hybridizes with nucleic acid encoding MUM-2. In an embodiment, the nucleic acid probe is complementary to nucleic acid encoding MUM-2.

In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-1 is a DNA probe. In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-2 is a DNA probe.

In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-1 is a RNA probe. In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-2 is a RNA probe.

In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-1 is a genomic DNA probe. In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-2 is a genomic DNA probe.

In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-1 is labeled with

a detectable marker. In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-2 is labeled with a detectable marker.

5 In an embodiment, the detectable marker is selected from the group consisting of a radioactive isotope, enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.

10 In an embodiment, the nucleic acid probe which specifically hybridizes with nucleic acid encoding MUM-1 is linked to a nucleic acid sequence capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule of human chromosome 14. In an embodiment, the nucleic acid probe which specifically
15 hybridizes with nucleic acid encoding MUM-2 is linked to a nucleic acid sequence capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule of human chromosome 14.

20 This invention provides a nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding a MUM protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid
25 molecule of human chromosome 14.

In an embodiment, the nucleic acid probe comprises a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid
30 molecule encoding a MUM-1 protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

In an embodiment, the nucleic acid probe comprises a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding a MUM-2 protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

In an embodiment, the nucleic acid probe comprises a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding a MUM-1 protein which is linked at a specific break point to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

In an embodiment, the nucleic acid probe comprises a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding a MUM-2 protein which is linked at a specific break point to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

In an embodiment, the specific break point of the nucleic acid probe comprises a portion of the t(6;14)(p25;q32) translocation. In an embodiment, the specific break point of the nucleic acid probe comprises a portion of a t(14;15) translocation. In an embodiment, the nucleic acid probe comprising a portion of the t(6;14)(p25;q32) translocation is labeled with a detectable marker. In an embodiment, the nucleic acid probe comprising a portion of a t(14;15) translocation is labeled with a detectable marker. In an

embodiment, the nucleic acid probe comprising a portion of the t(6;14)(p25;q32) or comprising a portion of a t(14;15) translocation of claim 60, has a detectable marker selected from the group consisting of a radioactive isotope, enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.

This invention provides a method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-1 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-1 protein.

This invention provides a method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-2 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-2 protein.

In an embodiment, the rearrangement of nucleic acid encoding MUM-1 protein is detected by contacting the nucleic acid from the sample with a MUM-1 probe under conditions permitting the MUM-1 probe to hybridize with the nucleic acid encoding MUM-1 protein from the sample, thereby detecting the rearrangement of nucleic acid encoding MUM-1 protein in the sample.

In an embodiment, the rearrangement of nucleic acid encoding MUM-2 protein is detected by contacting the nucleic acid from the sample with a MUM-2 probe under conditions permitting the MUM-2 probe to hybridize with the nucleic acid encoding MUM-2 protein from the sample, thereby

detecting the rearrangement of nucleic acid encoding MUM-2 protein in the sample.

5 In an embodiment, the rearrangement of nucleic acid encoding MUM-1 protein is detected by a MUM-1 probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-1 protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

15 In an embodiment, the rearrangement of nucleic acid encoding MUM-2 protein is detected by a the MUM-2 probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-2 protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 15.

20 In an embodiment, the MUM-1 probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-1 protein is linked at a specific break point to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

30 In an embodiment, the MUM-2 probe comprising a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-2 protein is linked at a specific break point to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 15.

35 In an embodiment, the MUM-1 probe comprises a specific break point comprising a portion of the t(6;14)(p25;q32)

translocation. In an embodiment, the MUM-2 probe comprises a specific break point comprising a portion of a t(14;15) translocation.

In an embodiment, the method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-1 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-1 protein comprises: a) obtaining DNA from the sample of the subject suffering from multiple myeloma; b) performing a restriction digest of the DNA with a panel of restriction enzymes; c) separating the resulting DNA fragments by size fractionation; d) contacting the resulting DNA fragments with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-1 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 and labeled with a detectable marker; e) detecting labeled bands which have hybridized to the nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-1 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 to create a unique band pattern specific to the DNA of subjects suffering from multiple myeloma; f) preparing DNA obtained from a sample of a subject for diagnosis by steps (a-e); and g) comparing the detected band pattern specific to the DNA obtained from a sample of subjects suffering from multiple myeloma from step (e) and the DNA obtained from a sample of the subject for diagnosis from step (f) to determine whether the patterns

are the same or different and to diagnose thereby predisposition to multiple myeloma if the patterns are the same.

5 In an embodiment, the method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-2 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-2 protein
10 comprises: a) obtaining DNA from the sample of the subject suffering from multiple myeloma; b) performing a restriction digest of the DNA with a panel of restriction enzymes; c) separating the resulting DNA fragments by size fractionation; d) contacting the resulting DNA fragments
15 with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-2 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 and labeled with a detectable marker; e) detecting labeled bands which have hybridized to the nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule
25 encoding a human MUM-2 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 to create a unique band pattern specific to the DNA of subjects suffering from multiple myeloma; f) preparing DNA obtained from a sample of a subject for diagnosis by steps (a-e); and g) comparing the detected band pattern specific to the DNA obtained from a sample of subjects suffering from multiple myeloma from step (e) and the DNA obtained from a sample of the subject for
30

diagnosis from step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to multiple myeloma if the patterns are the same.

5

In an embodiment, the size fractionation in step (c) is effected by a polyacrylamide or agarose gel. In an embodiment, the detectable marker is radioactive isotope, enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.

10

In an embodiment, the method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-1 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-1 protein comprises: a) obtaining RNA from the sample of the subject suffering from multiple myeloma; b) separating the RNA sample by size fractionation; c) contacting the resulting RNA species with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-1 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 and labeled with a detectable marker; d) detecting labeled bands which have hybridized to the RNA species to create a unique band pattern specific to the RNA of subjects suffering from multiple myeloma; e) preparing RNA obtained from a sample of a subject for diagnosis by steps (a-d); and f) comparing the detected band pattern specific to the RNA obtained from a sample of subjects suffering from multiple myeloma from step (d) and the RNA obtained from a sample of the subject for diagnosis from

15

20

25

30

35

step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to multiple myeloma if the patterns are the same.

5 In an embodiment, the method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-2 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-2 protein
10 comprises: a) obtaining RNA from the sample of the subject suffering from multiple myeloma; b) separating the RNA sample by size fractionation; c) contacting the resulting RNA species with a nucleic acid probe capable of specifically hybridizing with a unique sequence included
15 within the sequence of a nucleic acid molecule encoding a human MUM-2 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 15 and labeled with a detectable marker; d)
20 detecting labeled bands which have hybridized to the RNA species to create a unique band pattern specific to the RNA of subjects suffering from multiple myeloma; e) preparing RNA obtained from a sample of a subject for diagnosis by steps (a-d); and f) comparing the detected band pattern
25 specific to the RNA obtained from a sample of subjects suffering from multiple myeloma from step (d) and the RNA obtained from a sample of the subject for diagnosis from step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to multiple
30 myeloma if the patterns are the same.

In an embodiment, the size fractionation in step (b) is effected by a polyacrylamide or agarose gel. In an embodiment, the detectable marker is radioactive isotope,

enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.

5 In an embodiment, multiple myeloma associated with the expression of a specific human MUM-1 is diagnosed by the method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-1 protein in a DNA or RNA sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid
10 encoding MUM-1 protein.

15 In an embodiment, multiple myeloma associated with the expression of a specific human MUM-2 is diagnosed by the method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-12 protein in a DNA or RNA sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-2 protein.

20 This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to an mRNA molecule encoding a human MUM-1 protein so as to prevent overexpression of the mRNA molecule. This invention provides an antisense oligonucleotide having a sequence
25 capable of specifically hybridizing to an mRNA molecule encoding a human MUM-2 protein so as to prevent overexpression of the mRNA molecule.

30 This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to the cDNA molecule encoding a MUM protein. This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to the genomic DNA molecule

encoding a MUM protein. This invention provides an antisense oligonucleotide having a sequence capable of specifically hybridizing to the RNA molecule encoding a MUM protein.

5 This invention provides a purified MUM protein. This invention provides a purified MUM-1 protein. This invention provides a purified human MUM-1 protein. This invention provides an antibody directed to a purified MUM-1 protein. This invention provides an antibody capable of specifically
10 recognizing MUM-1 protein. In an embodiment, the antibody capable of specifically recognizing MUM-1 protein is a human MUM-1 protein.

This invention provides a purified MUM-2 protein. This
15 invention provides a purified human MUM-2 protein. This invention provides an antibody directed to a purified MUM-2 protein. This invention provides an antibody capable of specifically recognizing MUM-2 protein. In an embodiment, the antibody capable of specifically recognizing MUM-2
20 protein is a human MUM-2 protein.

In an embodiment, the antibody directed to a purified MUM-1 protein is a monoclonal antibody. In an embodiment, the antibody capable of specifically recognizing MUM-1 protein
25 is a monoclonal antibody. In an embodiment, the antibody capable of specifically recognizing MUM-1 protein is a human MUM-1 protein.

In an embodiment, the antibody directed to a purified MUM-2
30 protein is a monoclonal antibody. In an embodiment, the antibody capable of specifically recognizing MUM-2 protein is a monoclonal antibody. In an embodiment, the antibody capable of specifically recognizing MUM-2 protein is a human

MUM-2 protein.

5 This invention provides a pharmaceutical composition
comprising an amount of the oligonucleotide having a
sequence capable of specifically hybridizing to an mRNA
molecule encoding a human MUM-1 protein so as to prevent
overexpression of the mRNA molecule effective to prevent
overexpression of a human MUM-1 protein and a
pharmaceutically acceptable carrier capable of passing
10 through a cell membrane.

15 This invention provides a pharmaceutical composition
comprising an amount of the oligonucleotide having a
sequence capable of specifically hybridizing to a cDNA
molecule encoding a MUM protein effective to prevent
overexpression of a human MUM-1 protein and a
pharmaceutically acceptable carrier capable of passing
through a cell membrane.

20 This invention provides a pharmaceutical composition
comprising an amount of the oligonucleotide having a
sequence capable of specifically hybridizing to a genomic
DNA molecule effective to prevent overexpression of a human
MUM-1 protein and a pharmaceutically acceptable carrier
25 capable of passing through a cell membrane.

30 This invention provides a pharmaceutical composition
comprising an amount of the oligonucleotide having a
sequence capable of specifically hybridizing to an mRNA
molecule encoding a human MUM-2 protein so as to prevent
overexpression of the mRNA molecule effective to prevent
overexpression of a human MUM-2 protein and a
pharmaceutically acceptable carrier capable of passing

through a cell membrane.

5 This invention provides a pharmaceutical composition comprising an amount of the oligonucleotide having a sequence capable of specifically hybridizing to a cDNA molecule encoding a MUM protein effective to prevent overexpression of a human MUM-2 protein and a pharmaceutically acceptable carrier capable of passing through a cell membrane.

10

This invention provides a pharmaceutical composition comprising an amount of the oligonucleotide having a sequence capable of specifically hybridizing to a genomic DNA molecule effective to prevent overexpression of a human MUM-2 protein and a pharmaceutically acceptable carrier capable of passing through a cell membrane.

15

This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described more fully in the claims which follow thereafter.

20

25

EXPERIMENTAL DETAILS

Materials and Methods

5 **Cell lines.** The following myeloma cell lines were used in
the present study: SK-MM-1, RPMI-8226, U266, EJM, XG-1,
XG-2, XG-4, XG-5, XG-6, XG-7, and XG-10. The RPMI-8226 cell
line was obtained through the American Type Culture
Collection (ATCC, Rockville, MD). SKMM-1 and U-266 cell
10 lines were gifts from Dr. A. N. Houghton and Dr. K. Nilsson,
respectively (18; 12). Characterization of these cell lines
were previously reported. Six XG cell lines were gifts from
Dr. B. Klein and were cultured in RPMI 1640 containing 10%
fetal calf serum (FCS), 5×10^{-5} mol/L 2-ME, and rIL-
15 6 (1ng/mL) (13;19). Other myeloma cell lines used were all
IL-6 independent. The SK-MM-1 cell line was used to isolate
the chromosomal breakpoint carrying the 14q+ chromosome
without any information on the donor chromosome. XG-1,
XG-2, XG-6, XG-8 cell lines are reported to carry the
20 t(11;14)(q13;q32) translocation. XG-5 cells also share
both t(11;14) and t(8;14)(q24;q32).

Southern and Northern blot analyses. Southern blot analysis
was performed as previously described (21). Briefly, ten
25 micrograms of high molecular-weight DNA extracted from each
cell line was digested to completion with *Bam*HI and *Hind*III
restriction enzymes, size- fractionated on 0.7% agarose gel,
and transferred onto Duralose nitrocellulose membrane
(Stratagene) according to the manufacturer's instructions.
30 Blots were hybridized with a random-primed DNA probe and
washed at 60°C in 0.2 x SSC and 0.1 % SDS for 5 minutes.
Genomic probes used in this study were as follows; human IgH

J region JH probe (6.6kb *Bam*HI-*Hind*III fragment) was provided by Dr. J. V. Ravetch, human IgH C μ probe (1.3 kb *Eco*RI fragment) was provided by Dr. S.J. Korsmeyer. Human IgH region Cy2 probe was provided by Dr. C. Croce.

5

Northern blot analysis was performed as described previously (21). Briefly, a 10 μ g aliquot of total RNA was loaded on each lane and probed with a 2.1H probe of the MUM1 gene (Figure 2A). GAPDH or β -actin probes were used as controls for amount of total RNA.

10

Genomic library. High molecular-weight DNA of SK-MM-1 cell line was digested completely with *Bam*HI and partially with *Sau*3AI, and size-fractionated by using a low-melting point agarose gel. DNA ranging from 10kb to 23kb were purified and ligated into the *Bam*HI sites of λ -DASH II phage vector (Stratagene, La Jolla, CA). After packaging, 3×10^5 and 6×10^5 recombinant clones of the *Bam*HI digested library and partially digested library were screened with JH and C μ probes, respectively. To isolate the germline region of the 6p25 locus, a commercially available human placental library (Stratagene) was screened. Positive clones were mapped with restriction enzymes by partial digestion of the phage DNAs followed by probing with T7 and T3 primers labeled with T4 polynucleokinase and 32p- γ ATP.

15

20

25

cDNA library. A phage library constructed by oligo-dT and random-priming normal human spleen RNA (Clontech) was screened by 2.1H probe (Figure 2A) to isolate initial MUM1 cDNA clones. After the first round of screening, positive clones were used as probes to walk to the 5' side using the same library. Positive clones were subcloned into

30

pBluescript and analyzed for mapping and sequencing.

DNA sequencing. DNA sequences were determined by the dideoxy chain termination method and analyzed by an
5 ABI (Applied Biosystems) autosequencer. Deletion mutants for sequencing were prepared using exonuclease III and mung bean nuclease. cDNA sequences were analyzed with the Genetics Computer Group (GCG) programs. Sequence homology searches
10 were carried out through the BLAST E-mail server at the National Center for Biotechnology Information, National Library of Medicine, Bethesda, MD.

Fluorescence in situ hybridization (FISH). Metaphase chromosome from human lymphocytes were prepared. A
15 biotin-labeled probe was prepared by nick-translation using Bio-16-dUTP. Conditions for hybridization and washing were described previously (22).

Experimental Results

20 **IgH gene rearrangement of the SK-MM-I cell line.** In *Bam*HI digestion, the JH probe detects two rearranged bands of the size of 12.0 kb and 9.7kb (Fig 1). The 9.7 kb band is comigrated with that probed with C γ 2 probe, suggesting it to
25 be a physiological rearrangement, although this cell line secretes only λ chain. One allele of the C μ locus is deleted and another is rearranged (6.5 kb) without being comigrated with rearranged bands of JH. Hybridization with a C α probe showed only the germline band (data not shown).
30 These results suggested the possibility of the chromosomal breakpoint between JH and C μ locus. Hence, the 12.0 kb and the 6.5 kb bands detected by JH and C μ were considered to

represent unknown derivative chromosome and derivative 14 chromosome, respectively.

Molecular cloning of the t(6;14)(p25;q32) breakpoint.

A genomic library constructed with *Bam*HI complete digestion was screened with a JH probe to isolate the 12.0 kb *Bam*HI band. Another library constructed with *Sau*3AI partial digestion was screened with a C μ probe to isolate phage clones containing the 6.5 kb *Bam*HI fragment. Two phage clones, λ SKB-4a and λ SKS-3, considered to represent the unknown derivative and derivative 14 chromosomes respectively, were obtained (Fig 2A). A 0.7 kb *Bam*HI-*Hind*III probe (0.7B/H) of the λ SKS-3 was used to confirm the comigration with the rearranged 6.5 kb C μ band by Southern analysis (Fig 1). The chromosomal origin of the centromeric side of the λ SKB-4a and telomeric side of the λ SKS-3 were confirmed by hybridization to a somatic cell hybrid DNA panel with a 4.5 kb *Apa*I fragment (4.5A) and 2.1 kb *Hind*III (2.1H) probes. Both probes showed positive signals in hybrid cell DNA containing a human chromosome 6 (data not shown). These probes were also used to isolate the germline chromosome 6 region by screening the human placental genomic library. One of the phage clone DNA (λ MUM-3) was used as a probe for FISH analysis. It identified the localization of this region to be chromosome 6 short arm p25 (Fig 3). To investigate the precise breakpoint within the IgH gene, a 1.5 kb *Hind*III-*Eco*RI fragment of the λ SKS-3, containing the breakpoint on derivative 14 chromosome was sequenced. The breakpoint was confirmed to be just 3' to the switch μ (S μ) repetitive sequences (Fig 2B). Nucleotide sequencing of the region around the breakpoints of chromosome 6 and derivative 6

chromosome showed that the chromosomal translocation was reciprocal with minimum deletion of both the IgH and 6p25 sequences.

5 **Transcriptional unit in the vicinity of the 6p25 breakpoint.**

10 An attempt to find a functional transcriptional unit in the vicinity of the breakpoints was made. Although a 4.5A probe on derivative 6 chromosome could not detect any transcripts, a 2.1H probe on derivative 14 chromosome detected a single
15 6 kb transcript in the SK-MM-1 cell line. Accordingly, this gene was designated as *MUM1* (multiple myeloma oncogene 1). The same probe was used to study the expression of the *MUM1* gene in various hematopoietic cell lines. The 6 kb message was expressed at high levels in most B cell lines and at low
20 levels in peripheral T cell lines (Fig 4A). Cell lines derived from immature T cells, the myelomonocytic lineage, and erythroid lineage do not seem to express *MUM1*. In B cells, *MUM1* appears to be expressed throughout the development from the preB cell stage to the plasma cell stage (Fig 4B). However, some of the Burkitt's lymphoma derived cell lines such as BJA-B did not express this gene (data not shown). The expression level of the *MUM1* transcript in myeloma cell lines was also examined (Fig 4C).
25 The SK-MM-1 cell line showed a 7.5-fold overexpression when compared with the other three IL-6 independent cell lines, suggesting a deregulated expression of the translocated allele. It is of interest that the IL-6 dependent XG-4, XG-7, and XG-10 cell lines are also expressing at high
30 levels. Particularly, expression in the XG-7 cell line is 19.9 times the average of the aforementioned control cell lines.

MUMI cDNA cloning, sequencing, and homology search.

Human spleen cDNA library was initially screened with a 2.1H probe followed by three times walking to 5' side using cDNA probes. A 5.5kb cDNA, approximately corresponding to the size detected by Northern analysis was isolated. This cDNA contained a 1,353 base pair open reading frame (ORF) and a long 3' untranslated region (Fig 5A). The ORF encodes for a protein of 451 amino acids with a predicted molecular weight of 50 kD (Fig 5B). The putative ATG initiation codon at position 217 has G at the -3 position which corresponds to the Kozak consensus sequence (23). The ORF is preceded by two in-frame stop codons. A database search demonstrated a significant similarity between MUM-1 ORF and the interferon regulatory factor (IRF) family proteins. The NH₂-terminal of the MUM-1 ORF shares a high homology with all of the IRF family proteins which share a characteristic DNA binding motif consisting of the conserved 5 tryptophan residues (Fig 6A). The COOH-terminal also has a high homology with ICSBP (interferon consensus sequence binding protein)(21), ISGF3 γ (interferon-stimulated gene factor-3 gamma)(22), and IRF-3 protein (23)(Fig 6B), although it did not have any homologous regions with IRF-1 and IRF-2 protein. The highest similarity (95.1%) and identity (91.8%) were found with a possible mouse homolog, LSIRF (lymphoid specific interferon regulatory factor)/Pip (PU-1 cofactor protein-1)(24,25). A high similarity was found with ICSBP (63.98%), ISGF3 γ (55.8%), and IRF3 (50.1%) among the human IRF family protein members. A gene sequence encoding a nearly identical protein was recently deposited in GenBank. This gene, termed ICSAT (interferon consensus sequence binding protein in adult T-cell leukemia cell lines or activated T cells) is likely to be the same gene as MUM1

(26).

Breakpoints at MUM1 locus in multiple myeloma.

In order to analyze the exact location of the SK-MM-1
5 breakpoint at the 6p25 locus and to explore the frequency of
the MUM1 gene involvement in myeloma cases, we walked nearly
55kb in a human placental genomic phage library around the
MUM1 gene and determined the rough exon-intron structure as
shown in Figure 7 (Fig. 7). The SK-MM-1 breakpoint was
10 located 3' to the last exon, containing a poly A additional
signal, consistent with an unaltered size of the MUM1
transcript of this cell line in Northern analysis. Seven
repeat-free genomic probes shown in Figure 7 have been used
to investigate the rearrangement in Southern analyses of the
15 11 MM cell lines and 18 MM cases. One case (case 10)
displayed rearranged bands in *Bam*HI and *Xba*I digests when
analyzed using a 0.9A probe located at 3' to the MUM1 gene.

**Cloning of the MUM2 locus from the U-266 multiple myeloma
20 cell line.**

Using an experimental strategy analogous to the one
described for the cloning of the MUM1 gene from the SK-MM-1
cell line, a second genetic locus altered in multiple
myeloma (MUM2) was identified by analyzing the U-266
multiple myeloma cell line. Briefly, Southern blot analysis
25 using *Bam*H restriction digestion and various Ig probes showed
that U-266 DNA contained two rearranged fragments (shown by
arrowheads in Fig. 9) containing C α sequences and lacking J
sequences. These two fragments (der 14 and 14q32 in Fig. 9)
30 were cloned from a genomic library constructed from U-266
DNA along with a normal 14q 32 locus (14q32 germline in Fig.
9). In order to determine whether a gene was located in

proximity to the chromosomal breakpoints in der 14, the 2.5 BE restriction fragment (see Fig. 9), which was at the opposite side of the Ig Ca sequences, was used to probe a Northern blot carrying RNA from various MM cell lines. The results (Fig. 10) showed that a 1.9 kb mRNA was detectable in some of these cell lines including U-266. This result showed that a gene, called MUM2, normally not present within the Ig locus on chromosome 14q32, had been translocated in proximity of the Ig locus in U-266 cells. Since the Ig locus contains strong transcriptional regulatory elements, it is likely that the expression of this gene is deregulated in these cells. The structure of the MUM2 gene and its protein are currently under investigation. The 2.5 BE probe and other probes derived from the der 14 phage can be used to screen MM cases for MUM2 rearrangements as shown for MUM1 (Fig. 7).

Experimental Discussion

Using the experimental strategies used for the identification of the MUM1 and MUM2 genes in the SK-MM-1 and U-266 cell lines, respectively, it is possible to analyze most MM cases and isolate the corresponding genes. The scheme shown in Fig. 11 shows that the physiological IgH gene rearrangements (Fig. 11A) typically maintain linkage of C and J sequences and this linkage becomes detectable by using an appropriate restriction enzyme digestion (BamHI in the example in Fig. 11). Conversely, chromosomal translocations (14q+) affecting the IgH locus on 14q32 lead to breakage of the C-J linkage and the two sets of sequences appear on distinct restriction fragments. (Fig. 11B) Table 1 shows the application of this analysis to a panel of MM

cell lines and biopsies. The results show that at least 65% of cases show breakage of the C-J linkage within Ig J or switch regions. The restriction fragments containing either C or J sequences (R in Table 1) can be cloned as shown for the SK-MM-1 and U-266 cell lines and the genes flanking the chromosomal breakpoints can be used as probes to screen additional MM cases for similar rearrangements, whereas the sequence of the genes can be used to understand the consequences of these genetic lesions in multiple myeloma. Cloning of the chromosomal breakpoints and corresponding genes is currently ongoing for all of the MM cases shown in Table 1.

The method of analysis of 14q+ chromosomal translocations and identification of the genes altered in multiple myeloma of this invention will allow 1) the determination of chromosomal sequences involved in 14q+ translocations, the most important cytogenetic lesion associated with MM pathogenesis elucidation; 2) elucidation of specific gene lesions for MM; 3) a diagnostic method based on gene/DNA lesion and 4) a therapeutic approach aimed at counteracting the action of abnormal gene products.

Table 1. Summary of JH-C breakage analysis in MM cell lines and biopsies (cases). Rearrangement (R) involving physiologic Ig recombinations, i.e. retaining JH-C linkage are marked as R*; rearrangements lacking JH-C linkage, and therefore suggesting a 14q+ chromosomal breakpoint, are marked as R. The latter represents candidates for cloning an further analysis.

Cell Line/Case	sIg	JH	Cμ	Cα	Sγ3'	possible breakpoint locus
10 RPMI-8226	λ	D/D	D/D	G	<u>R</u> /G	Sγ
U-266	Eλ	R/D	D/D	<u>R</u> / <u>R</u> /G	G	Sα
EJM	Gλ	<u>R</u> /R*	D/D	G	R*/G	JH~Sμ
XG-1	Aκ	R*/D	D/D	R*	G	ND
XG-2	Gλ	R*/D	D/D	G	R*/G	ND
15 XG-4	Gκ	R*/ <u>R</u>	D/D	G	R*/G	JH~Sμ
XG-5	λ	<u>R</u> /D	D/D	G	G	JH~Sμ
XG-6	Gλ	R*/ <u>R</u>	D/D	G	R*/G	JH~Sμ
XG-7	Aκ	R/D	D/D	R/G	<u>R</u> /D	Sγ
XG-10	G	R*/ <u>R</u>	D/D	G	R*/G	JH~Sμ
20 SK-MM-1	κ	R*/ <u>R</u>	<u>R</u> /D	G	R*/G	JH~Sμ
CASE125		R*	G	G	R*	ND
CASE33		<u>R</u> /R*	G	G	<u>R</u> /R*	Sγ
CASE34		R*	G	R*	G	ND
25 CASE93		R*	G	R*	G	ND
CASE91		R*	R*	<u>R</u>	G	Sα
CASE128		R*	G	R*	G	ND

R*, comigrated bands with JH; R, target bands to isolate; ND, not determined

Possible breakage in switch regions:

Cell Lines	4/11 (36%)	
Cases	2/6 (33%)	Total 6/17 (35%)

Possible breakage in JH ~ switch regions:

Cell Lines	9/11 (82%)	
Cases	2/6 (33%)	Total 11/17 (65%)

References

1. Rabbitts, T.H. Chromosomal translocations in human cancer. Nature 372:143-149, 1994.
5
2. Dalla-Favera, R., Bregni, M., Erickson, D., Patterson, D., Gallo, R.C., Croce, C.M. Human c-myc oncogene is located on the region of chromosome 8 that is translocated in Burkitt lymphoma cells. Proc Nat Acad Sci USA 79:7824, 1982.
10
3. Tsujimoto, Y., Jaffe, E., Cossman, J., Gorham, J., Nowell, P.C., Croce, C.M. Clustering of breakpoints on chromosome 11 in human B-cell neoplasms with the t(11;14) chromosome translocation. Nature 315:340-343, 1985.
15
4. Motokura, T., Bloom, T., Kim, H.G., Juppner, H., Ruderman, J.V., Kronenberg, H.M., Arnold, A. A novel cyclin encoded by a bcl-1 linked candidate oncogene. Nature 350:512-515, 1991.
20
5. Tsujimoto, Y., Yunis, J., Onorato-Showe, L., Erikson, J., Nowell, P.C., Croce, C.M. Molecular cloning of chromosomal breakpoint of B-cell lymphomas and leukemias with the t(11;14) chromosome translocation. Science 224:1403-1406, 1984.
25
6. Cleary, M.L., Sklar, J. : Nucleotide sequence of a t(14;18) chromosomal breakpoint in follicular lymphoma and demonstration of a breakpoint-cluster region near a transcriptionally active locus on chromosome 18. Proc
30

Natl Acad Sci USA 82: 7439, 1985.

7. Bakhshi, A., Jensen, J.P., Goldman, P., Wright, J.J.,
5 McBride, O.W., Epstein, A.L., Korsmeyer, S.J. Cloning
the chromosomal breakpoint of t(14;18) human lymphomas:
clustering around JH on chromosome 14 and near a
transcriptional unit on 18. Cell 41:889, 1985.
8. Ye, B.H., Lista, F., Lo Coco, F., Knowles, D.M., Offit,
10 K., Chaganti, R.S.K., Dalla-Favera, R. Alterations of
a zinc finger-encoding gene, BCL-6, in diffuse
large-cell lymphoma. Science 262:747-750, 1993.
9. Dewald, G.W., Kyle, R.A., Hicks, G.A., Greipp, P.R. The
15 clinical significance of cytogenetic studies in 100
patients with multiple myeloma, plasma cell leukemia,
or amyloidosis. Blood 66:380-390, 1985.
10. Gould, J., Alexanian, R., Goodacre, A., Pathak, S.,
20 Hecht, B., Barlogie, B. Plasma cell karyotype in
multiple myeloma. Blood 71:453-456, 1988.
11. Weh, H.J., Gutensohn, K., Selbach, J., Kruse, R.,
Wacker-Backhaus, G., Seeger, D., Fiedler, W., Fett, W.,
25 Hossfeld, D.K. Karyotype in multiple myeloma and plasma
cell leukemia. Eur J Cancer 29A:1269-1273, 1993.
12. Jernberg, H., Zech, L., Nilsson, K. Cytogenetic studies
on human myeloma cell lines. Int J Cancer 40: 811-817,
30 1987.
13. Zhang, X-G., Gaillard, J.P., Robillard, N., Lu, Z-Y.,

Gu, Z-G., Jourdan M., Boiron, J.M., Bataille, R., Klein, B. Reproducible obtaining of human myeloma cell lines as a model for tumor stem cell study in human multiple myeloma. Blood 83:3654-3663, 1994.

5

14. Seto, M., Yamamoto, K., Iida, S., Akao, Y., Utsumi, K.R., Kubonishi, I., Miyoshi, I., Ohtsuki, T., Yawata, Y., Namba, M., Motokura, T., Arnold, A., Takahashi, T., Ueda, R. Gene rearrangement and overexpression of PRAD1 in lymphoid malignancy with t(11;14)(q13;q32) translocation. Oncogene 7:1401 -1406, 1992.

10

15. Rabbitts, P.H., Douglas, J., Fisher, P., Nacheva, E., Karpas, A., Catovsky, D., Melo, J.V., Baer, R., Stinson, M.A., Rabbitts, T.H. Chromosome abnormalities at 11q13 in B cell tumours. Oncogene 3:99103, 1988.

15

16. Fiedler, W., Weh H.J., Hossfeld, D.K. Comparison of chromosome analysis and BCL-1 rearrangement in a series of patients with multiple myeloma. Br J Haematol 81: 58-61, 1992.

20

17. Taniwaki, M., Nishida, K., Takashima, T., Nakagawa, H., Fujii, H., Tamaki, T., Shimazaki, C., Horiike, S., Misawa, S., Kashima, K. Nonrandom chromosomal rearrangements of 14q32.3 and 19p13.3 and preferential deletion of 1p in 21 patients with multiple myeloma and plasma cell leukemia. Blood 84: 2283-2290, 1994.

25

18. Sun, Z., Kitchingman, G.R. Sequencing of selected regions of the human immunoglobulin heavy-chain gene locus that completes the sequence from JH through delta

30

constant region. DNA sequence 1:347-355, 1991.

19. Eton, O., Scheinberg, D.A., Houghton, A.N.
Establishment and characterization of two human myeloma
cell lines secreting kappa light chains. *Leukemia* 3:
729-735, 1989.
20. Mazars, G-R., Portier, M., Zhang, X-G., Jourdan, M.,
Bataille, R., Theillet, C., Klein, B. Mutations of the
pS3 gene in human myeloma cell lines. *Oncogene* 7:
1015-1018, 1992.
21. Iida, S., Seto M., Yamamoto, K., Tojo, A., Asano, S.,
Kamada, N., Ariyoshi, Y., Takahashi, T., Ueda, R. MLLT3
gene on 9p22 involved in t(9;11) leukemia encodes a
serine/proline rich protein homologous to MLLT1 on
19p13. *Oncogene* 8(11):3085-3095, 1993.
22. Rao, P.H., Murty, V.V.V.S., Gaidano, G., Hauptschein,
R., Dalla-Favera, R., and Chaganti, R.S.K. Subregional
mapping of 8 single copy loci to chromosome 6 by
fluorescence in situ hybridization. *Cytogenet. Cell
Genet.* 66:272-273, 1994.
23. Kozak, M. The scanning model for translation: an
update. *J. Cell. Biol.* 108:229-241, 1989.
24. Driggers, P.H., Ennist, D.L., Gleason, S.L., Mak, W-H.,
Marka, M.S., Levi, B-Z., Flanagan, J.R., Appella, E.,
Ozato, K. : *Proc Natl Acad Sci USA* 87: 3743-3747, 1990.
25. Veals, S.A., Schindler, C., Leonardo, D., Fu, X-Y.,

- 5 Aebersold, R., Damell, J.E., Levy, D.E. Subunit of an
 alpha-interferon-responsive transcription factor is
 related to interferon regulatory factor and myb
 families of DNA-binding proteins. Mol Cell Biol 12:
 3315-3324, 1992.
- 10 26. Grant, C.E., Vasa, M.Z., Deeley, R.G. cIRF-3, a new
 member of the interferon regulatory factor(IRF) family
 that is rapidly and transiently induced by dsRNA.
 Nucleic Acid Res 23:2137-2145, 1995.
- 15 27. Matsuyama, T., Grossman, A., Mittrucker, H-W.,
 Siderovski, D.P., Kiefer, F., Kawakami, T., Richardson,
 C.D., Taniguchi, T., Yoshinaga, S.K., Mak, T.W.
 Molecular cloning of LSIRF, a lymphoid-specific member
 of the interferon regulatory factor family that binds
 the interferon-stimulated response element(ISRE).
 Nucleic Acid Res 23:2127-2136, 1995.
- 20 28. Eisenbeis, C.F., Singh, H., Storb, U. Pip, a novel IRF
 family member, is a lymphoid-specific, PU.1-dependent
 transcriptional activator. Genes & Dev 9:1377-1387,
 1995.
- 25 29. Yamagata, T., Nishida, J., Tanaka, T., Sakai, R.,
 Mitani, K., Yoshida, M., Taniguchi, T., Yazaki, Y.,
 Hirai, H. A novel interferon regulatory factor family
 transcription factor, ICSAT/Pip/LSIRF, that negatively
 regulates the activity of interferon-regulated genes.
30 Mol Cell Biol 16:1283-1294, 1996.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT: Dalla-Favera, Riccardo

(ii) TITLE OF INVENTION: IDENTIFICATION OF GENES ALTERED IN
MULTIPLE MYELOMA

10

(iii) NUMBER OF SEQUENCES: 17

(iv) CORRESPONDENCE ADDRESS:

15

(A) ADDRESSEE: Cooper & Dunham LLP

(B) STREET: 1185 Avenue of the Americas

(C) CITY: New York

(D) STATE: New York

(E) COUNTRY: U.S.A.

(F) ZIP: 10036

20

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

25

(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE: 28-MAY-1996

30

(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: White, John P.

(B) REGISTRATION NUMBER: 28,678

35

(C) REFERENCE/DOCKET NUMBER: 50995

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (212) 278-0400

(B) TELEFAX: (212) 391-0525

40

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 108 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Lys Leu Arg Gln Trp Leu Ile Asp Gln Ile Asp Ser Gly Lys Tyr Pro
1 5 10 15
Gly Leu Val Trp Glu Asn Glu Glu Lys Ser Ile Phe Arg Ile Pro Trp
20 25 30
Lys His Ala Gly Lys Gln Asp Tyr Asn Arg Glu Glu Asp Ala Ala Leu
35 40 45
Phe Lys Ala Trp Ala Leu Phe Lys Gly Lys Phe Arg Glu Gly Ile Asp
50 55 60
Lys Pro Asp Pro Pro Thr Trp Lys Thr Arg Leu Arg Cys Ala Leu Asn
65 70 75 80
Lys Ser Asn Asp Phe Glu Glu Leu Val Glu Arg Ser Gln Leu Asp Ile
85 90 95
Ser Asp Pro Tyr Lys Val Tyr Arg Ile Val Pro Glu
100 105

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 108 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

10 Lys Leu Arg Gln Trp Leu Ile Asp Gln Ile Asp Ser Gly Lys Tyr Pro
1 5 10 15

Gly Leu Val Trp Glu Asn Glu Glu Lys Ser Val Phe Arg Ile Pro Trp
20 25 30

15 Lys His Ala Gly Lys Gln Asp Tyr Asn Arg Glu Glu Asp Ala Ala Leu
35 40 45

Phe Lys Ala Trp Ala Leu Phe Lys Gly Lys Phe Arg Glu Gly Ile Asp
20 50 55 60

Lys Pro Asp Pro Pro Thr Trp Lys Thr Arg Leu Arg Cys Ala Leu Asn
65 70 75 80

25 Lys Ser Asn Asp Phe Glu Glu Leu Val Glu Arg Ser Gln Leu Asp Ile
85 90 95

Ser Asp Pro Tyr Lys Val Tyr Arg Ile Val Pro Glu
100 105

30

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 108 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: peptide

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

5 Arg Met Arg Pro Trp Leu Glu Met Gln Ile Asn Ser Asn Gln Ile Pro
1 5 10 15

Gly Leu Ile Trp Ile Asn Lys Glu Glu Met Ile Phe Gln Ile Pro Trp
20 25 30

10 Lys His Ala Ala Lys His Gly Trp Asp Ile Asn Lys Asp Ala Cys Leu
35 40 45

Phe Arg Ser Trp Ala Ile His Thr Gly Arg Tyr Lys Ala Gly Glu Lys
50 55 60

15 Glu Pro Asp Pro Lys Thr Trp Lys Ala Asn Phe Arg Cys Ala Met Asn
65 70 75 80

Ser Leu Pro Asp Ile Glu Glu Val Lys Asp Gln Lys Arg Asn Lys Gly
85 90 95

20 Ser Ser Ala Val Arg Val Tyr Arg Met Leu Pro Pro
100 105

(2) INFORMATION FOR SEQ ID NO:4:

25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 108 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
30 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

40 Arg Met Arg Pro Trp Leu Glu Glu Gln Ile Asn Ser Asn Thr Ile Pro
1 5 10 15

Gly Leu Lys Trp Leu Asn Lys Glu Lys Lys Ile Phe Gln Ile Pro Trp

20 25 30

Met His Ala Ala Arg His Gly Trp Asp Val Glu Lys Asp Ala Pro Leu
35 40 45

5

Phe Arg Asn Trp Ala Ile His Thr Gly Lys His Gln Pro Gly Val Asp
50 55 60

Lys Pro Asp Pro Lys Thr Trp Lys Ala Asn Phe Arg Cys Ala Met Asn
10 65 70 75 80

Ser Leu Pro Asp Ile Glu Glu Val Lys Asp Lys Ser Ile Lys Lys Gly
85 90 95

15

Asn Asn Ala Phe Arg Val Tyr Arg Met Leu Pro Leu
100 105

(2) INFORMATION FOR SEQ ID NO:5:

20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 107 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: peptide

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Arg Leu Arg Gln Trp Leu Ile Glu Gln Ile Asp Ser Ser Met Tyr Pro
1 5 10 15

35

Gly Leu Ile Trp Glu Asn Glu Glu Lys Ser Met Phe Arg Ile Pro Trp
20 25 30

Lys His Ala Gly Lys Gln Asp Tyr Asn Gln Glu Val Asp Ala Ser Ile
40 35 40 45

Phe Lys Ala Trp Ala Val Phe Lys Gly Lys Phe Lys Glu Gly Asp Lys

	50		55		60											
	Ala	Glu	Pro	Ala	Thr	Trp	Lys	Thr	Arg	Leu	Arg	Cys	Ala	Leu	Asn	Lys
	65					70					75					80
5																
	Ser	Pro	Asp	Phe	Glu	Glu	Val	Thr	Asp	Arg	Ser	Gln	Leu	Asp	Ile	Ser
					85					90					95	
	Glu	Pro	Tyr	Lys	Val	Tyr	Arg	Ile	Val	Pro	Glu					
10					100				105							

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 107 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: peptide

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

	Lys	Leu	Arg	Asn	Trp	Val	Val	Glu	Gln	Val	Glu	Ser	Gly	Gln	Phe	Pro
	1				5					10					15	
	Gly	Val	Cys	Trp	Asp	Asp	Thr	Ala	Lys	Thr	Met	Phe	Arg	Ile	Pro	Trp
					20				25					30		
	Lys	His	Ala	Gly	Lys	Gln	Asp	Phe	Arg	Glu	Asp	Gln	Asp	Ala	Ala	Phe
					35			40				45				
35																
	Phe	Lys	Ala	Trp	Ala	Ile	Phe	Lys	Gly	Lys	Tyr	Lys	Glu	Gly	Asp	Thr
		50					55					60				
	Gly	Gly	Pro	Ala	Val	Trp	Lys	Thr	Arg	Leu	Arg	Cys	Ala	Leu	Asn	Lys
40	65					70					75					80
	Ser	Ser	Glu	Phe	Lys	Glu	Val	Pro	Glu	Arg	Gly	Arg	Met	Asp	Val	Ala

85

90

95

Glu Pro Tyr Lys Val Tyr Gln Leu Leu Pro Pro
100 105

5

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 107 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

20

Arg Ile Leu Pro Trp Leu Val Ser Gln Leu Asp Leu Gly Gln Leu Glu
1 5 10 15

25

Gly Val Ala Trp Val Asn Lys Ser Arg Thr Arg Phe Arg Ile Pro Trp
20 25 30

Lys His Gly Leu Arg Gln Asp Ala Gln Gln Glu Asp Phe Gly Ile Phe
35 40 45

30

Gln Ala Trp Ala Glu Ala Thr Gly Ala Tyr Val Pro Gly Arg Asp Lys
50 55 60

35

Pro Asp Leu Pro Thr Trp Lys Arg Asn Phe Arg Ser Ser Ala Leu Asn
65 70 75 80

Arg Lys Glu Gly Leu Arg Leu Ala Glu Asp Arg Ser Lys Asp Pro His
85 90 95

40

Asp Pro His Lys Ile Tyr Glu Phe Val Asn Ser
100 105

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 95 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Lys	Arg	Leu	Cys	Gln	Ser	Thr	Ile	Tyr	Trp	Asp	Gly	Pro	Leu	Ala	Leu
1			5					10					15		
Cys	Asn	Asp	Arg	Pro	Asn	Lys	Leu	Glu	Arg	Asp	Gln	Thr	Cys	Lys	Leu
			20					25					30		
Phe	Asp	Thr	Gln	Gln	Phe	Leu	Ser	Glu	Leu	Gln	Ala	Phe	Ala	His	His
			35				40					45			
Gly	Arg	Ser	Leu	Pro	Arg	Phe	Gln	Val	Thr	Leu	Cys	Phe	Gly	Glu	Glu
			50				55				60				
Phe	Pro	Asp	Pro	Gln	Arg	Gln	Arg	Lys	Leu	Ile	Thr	Ala	His	Val	Glu
65				70					75						80
Pro	Leu	Leu	Ala	Arg	Gln	Leu	Tyr	Tyr	Phe	Ala	Gln	Gln	Asn	Ser	
				85					90					95	

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 95 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

5	Lys	Arg	Leu	Cys	Gln	Ser	Arg	Ile	Tyr	Trp	Asp	Gly	Pro	Leu	Ala	Leu
	1				5					10					15	
	Cys	Ser	Asp	Arg	Pro	Asn	Lys	Leu	Glu	Arg	Asp	Gln	Thr	Cys	Lys	Leu
				20					25					30		
10	Phe	Asp	Thr	Gln	Gln	Phe	Leu	Ser	Glu	Leu	Gln	Val	Phe	Ala	His	His
			35					40					45			
	Gly	Arg	Pro	Ala	Pro	Arg	Phe	Gln	Val	Thr	Leu	Cys	Phe	Gly	Glu	Glu
		50					55					60				
15	Phe	Pro	Asp	Pro	Gln	Arg	Gln	Arg	Lys	Leu	Ile	Thr	Ala	His	Val	Glu
	65					70					75					80
	Pro	Leu	Leu	Ala	Arg	Gln	Leu	Tyr	Tyr	Phe	Ala	Gln	Gln	Asn	Thr	
20					85					90					95	

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 96 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: peptide

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

	Lys	Arg	Leu	Cys	Gln	Gly	Arg	Val	Phe	Cys	Ser	Gly	Asn	Ala	Val	Val
	1				5					10					15	
40	Cys	Lys	Gly	Arg	Pro	Asn	Lys	Leu	Glu	Arg	Asp	Glu	Val	Val	Gln	Val
				20					25					30		

Phe Asp Thr Ser Gln Phe Phe Arg Glu Leu Gln Gln Phe Tyr Asn Ser
35 40 45

5 Gln Gly Arg Leu Pro Asp Gly Arg Val Val Leu Cys Phe Gly Glu Glu
50 55 60

Phe Pro Asp Met Ala Pro Leu Arg Ser Lys Leu Ile Leu Val Gln Ile
65 70 75 80

10 Glu Gln Leu Tyr Val Arg Gln Leu Ala Glu Glu Ala Gly Lys Ser Cys
85 90 95

(2) INFORMATION FOR SEQ ID NO:11:

15

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 96 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

20

- (ii) MOLECULE TYPE: peptide

25

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

30

Gln Arg Leu Cys Pro Ile Pro Ile Ser Trp Asn Ala Pro Gln Ala Pro
1 5 10 15

Pro Gly Pro Gly Pro His Leu Leu Pro Ser Asn Glu Cys Val Glu Leu
20 25 30

35

Phe Arg Thr Ala Tyr Phe Cys Arg Asp Leu Val Arg Tyr Phe Gln Gly
35 40 45

Leu Gly Pro Pro Pro Lys Phe Gln Val Thr Leu Asn Phe Trp Glu Glu
50 55 60

40

Ser His Gly Ser Ser His Thr Pro Gln Asn Leu Ile Thr Val Lys Met
65 70 75 80

Glu Gln Ala Phe Ala Arg Tyr Leu Leu Glu Gln Thr Pro Glu Gln Gln
85 90 95

5 (2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 100 amino acids

(B) TYPE: amino acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

20 Gln Arg Leu Gly His Cys His Thr Tyr Trp Ala Val Ser Glu Glu Leu
1 5 10 15

Leu Pro Asn Ser Gly His Gly Pro Asp Gly Glu Val Pro Lys Asp Lys
20 25 30

25

Glu Gly Gly Val Phe Asp Leu Gly Pro Phe Ile Val Asp Leu Ile Thr
35 40 45

30

Phe Thr Glu Gly Ser Gly Arg Ser Pro Arg Tyr Ala Trp Leu Phe Cys
50 55 60

Val Gly Glu Ser Trp Pro Gln Asp Gln Pro Trp Thr Lys Arg Leu Val
65 70 75 80

35

Met Val Lys Val Val Pro Thr Cys Leu Arg Ala Leu Val Glu Met Ala
85 90 95

Arg Val Gly Gly
100

40

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5176 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: other nucleic acid

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 217..1569

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

15

GCCTGACCAA CATGGTAAAA CCCCATCTCT GCTAAACTA CAAAAATTA GCTGGATGTG
60

20

GTGGCAGGGA ACCTGTCATC CCAGCTAGTT GGGAGACTGA GGCAGGAGAA TCGCTCGATC
120

TTGGGACCCA CCGCTGCCCT CAGCTCCGAG TCCAGGGCGA GTGCAGAGCA CAGCGGGCGG
180

25

AGGACCCCGG GCGCGGGCGC GGACGGCACG CGGGGC ATG AAC CTG GAG GGC GGC
234

Met Asn Leu Glu Gly Gly
1 5

30

GGC CGA GGC GGA GAG TTC GGC ATG AGC GCG GTG AGC TGC GGC AAC GGC
282

Gly Arg Gly Gly Glu Phe Gly Met Ser Ala Val Ser Cys Gly Asn Gly
10 15 20

35

AAG CTC CGC CAG TGG CTG ATC GAC CAG ATC GAC AGC GGC AAG TAC CCC
330

Lys Leu Arg Gln Trp Leu Ile Asp Gln Ile Asp Ser Gly Lys Tyr Pro
25 30 35

40

GGG CTG GTG TGG GAG AAC GAG GAG AAG AGC ATC TTC CGC ATC CCC TGG
378

Gly Leu Val Trp Glu Asn Glu Glu Lys Ser Ile Phe Arg Ile Pro Trp
40 45 50

5 AAG CAC GCG GGC AAG CAG GAC TAC AAC CGC GAG GAG GAC GCC GCG CTC
426
Lys His Ala Gly Lys Gln Asp Tyr Asn Arg Glu Glu Asp Ala Ala Leu
55 60 65 70

10 TTC AAG GCT TGG GCA CTG TTT AAA GGA AAG TTC CGA GAA GGC ATC GAC
474
Phe Lys Ala Trp Ala Leu Phe Lys Gly Lys Phe Arg Glu Gly Ile Asp
75 80 85

15 AAG CCG GAC CCT CCC ACC TGG AAG ACG CGC CTG CGG TGC GCT TTG AAC
522
Lys Pro Asp Pro Pro Thr Trp Lys Thr Arg Leu Arg Cys Ala Leu Asn
90 95 100

20 AAG AGC AAT GAC TTT GAG GAA CTG GTT GAG CGG AGC CAG CTG GAC ATC
570
Lys Ser Asn Asp Phe Glu Glu Leu Val Glu Arg Ser Gln Leu Asp Ile
105 110 115

25 TCA GAC CCG TAC AAA GTG TAC AGG ATT GTT CCT GAG GGA GCC AAA AAA
618
Ser Asp Pro Tyr Lys Val Tyr Arg Ile Val Pro Glu Gly Ala Lys Lys
120 125 130

30 GGA GCC AAG CAG CTC ACC CTG GAG GAC CCG CAG ATG TCC ATG AGC CAC
666
Gly Ala Lys Gln Leu Thr Leu Glu Asp Pro Gln Met Ser Met Ser His
135 140 145 150

35 CCC TAC ACC ATG ACA ACG CCT TAC CCT TCG CTC CCA GCC CAG CAG GTT
714
Pro Tyr Thr Met Thr Thr Pro Tyr Pro Ser Leu Pro Ala Gln Gln Val
155 160 165

40 CAC AAC TAC ATG ATG CCA CCC CTC GAC CGA AGC TGG AGG GAC TAC GTC
762
His Asn Tyr Met Met Pro Pro Leu Asp Arg Ser Trp Arg Asp Tyr Val
170 175 180

CCG GAT CAG CCA CAC CCG GAA ATC CCG TAC CAA TGT CCC ATG ACG TTT
810

Pro Asp Gln Pro His Pro Glu Ile Pro Tyr Gln Cys Pro Met Thr Phe
185 190 195

5

GGA CCC CGC GGC CAC CAC TGG CAA GGC CCA GCT TGT GAA AAT GGT TGC
858

Gly Pro Arg Gly His His Trp Gln Gly Pro Ala Cys Glu Asn Gly Cys
200 205 210

10

CAG GTG ACA GGA ACC TTT TAT GCT TGT GCC CCA CCT GAG TCC CAG GCT
906

Gln Val Thr Gly Thr Phe Tyr Ala Cys Ala Pro Pro Glu Ser Gln Ala
215 220 225 230

15

CCC GGA GTC CCC ACA GAG CCA AGC ATA AGG TCT GCC GAA GCC TTG GCG
954

Pro Gly Val Pro Thr Glu Pro Ser Ile Arg Ser Ala Glu Ala Leu Ala
235 240 245

20

TTC TCA GAC TGC CGG CTG CAC ATC TGC CTG TAC TAC CGG GAA ATC CTC
1002

Phe Ser Asp Cys Arg Leu His Ile Cys Leu Tyr Tyr Arg Glu Ile Leu
250 255 260

25

GTG AAG GAG CTG ACC ACG TCC AGC CCC GAG GGC TGC CGG ATC TCC CAT
1050

Val Lys Glu Leu Thr Thr Ser Ser Pro Glu Gly Cys Arg Ile Ser His
265 270 275

30

GGA CAT ACG TAT GAC GCC AGC AAC CTG GAC CAG GTC CTG TTC CCC TAC
1098

Gly His Thr Tyr Asp Ala Ser Asn Leu Asp Gln Val Leu Phe Pro Tyr
280 285 290

35

CCA GAG GAC AAT GGC CAC AGG AAA AAC ATT GAG AAC CTG CTG AGC CAC
1146

Pro Glu Asp Asn Gly His Arg Lys Asn Ile Glu Asn Leu Leu Ser His
295 300 305 310

40

CTG GAG AGG GGC GTG GTC CTC TGG ATG GCC CCC GAC GGG CTC TAT GCG
1194

Leu Glu Arg Gly Val Val Leu Trp Met Ala Pro Asp Gly Leu Tyr Ala
315 320 325

5 AAA AGA CTG TGC CAG AGC ACG ATC TAC TGG GAC GGG CCC CTG GCG CTG
1242

Lys Arg Leu Cys Gln Ser Thr Ile Tyr Trp Asp Gly Pro Leu Ala Leu
330 335 340

10 TGC AAC GAC CGG CCC AAC AAA CTG GAG AGA GAC CAG ACC TGC AAG CTC
1290

Cys Asn Asp Arg Pro Asn Lys Leu Glu Arg Asp Gln Thr Cys Lys Leu
345 350 355

15 TTT GAC ACA CAG CAG TTC TTG TCA GAG CTG CAA GCG TTT GCT CAC CAC
1338

Phe Asp Thr Gln Gln Phe Leu Ser Glu Leu Gln Ala Phe Ala His His
360 365 370

20 GGC CGC TCC CTG CCA AGA TTC CAG GTG ACT CTA TGC TTT GGA GAG GAG
1386

Gly Arg Ser Leu Pro Arg Phe Gln Val Thr Leu Cys Phe Gly Glu Glu
375 380 385 390

25 TTT CCA GAC CCT CAG AGG CAA AGA AAG CTC ATC ACA GCT CAC GTA GAA
1434

Phe Pro Asp Pro Gln Arg Gln Arg Lys Leu Ile Thr Ala His Val Glu
395 400 405

30 CCT CTG CTA GCC AGA CAA CTA TAT TAT TTT GCT CAA CAA AAC AGT GGA
1482

Pro Leu Leu Ala Arg Gln Leu Tyr Tyr Phe Ala Gln Gln Asn Ser Gly
410 415 420

35 CAT TTC CTG AGG GGC TAC GAT TTA CCA GAA CAC ATC AGC AAT CCA GAA
1530

His Phe Leu Arg Gly Tyr Asp Leu Pro Glu His Ile Ser Asn Pro Glu
425 430 435

40 GAT TAC CAC AGA TCT ATC CGC CAT TCC TCT ATT CAA GAA TGAAAAATGT
1579

Asp Tyr His Arg Ser Ile Arg His Ser Ser Ile Gln Glu
440 445 450

CAAGATGAGT GGT TTTCTTTT TTCCTTTTTT TTTT TTTT TTTTGATACG GAGATACGGG
1639

5 GTCTTGCTCT GTCTCCCAGG CTGGAGTGCA GTGACACAAT CTCAGCTCAC TGTGACCTCC
1699

GCCTCCTGGG TTCAAGAGAC TCTCCTGCCT CAGCCTCCCT GGTAGCTGGG ATTACAGGTG
1759

10 TGAGCCACTG CACCCACCCA AGACAAGTGA TTTTCATTGT AAATATTTGA CTTTAGTGAA
1819

AGCGTCCAAT TGA CTGCCCT CTTACTGTTT TGAGGAACTC AGAAGTGGAG ATTTCA GTTC
1879

15 AGCGGTTGAG GAGAATTGCG GCGAGACAAG CATGGAAAAT CAGTGACATC TGATTGGCAG
1939

20 ATGAGCTTAT TTCAAAGGA AGGGTGGCTT TGCATTTTCT TGTGTTCTGT AGACTGCCAT
1999

CATTGATGAT CACTGTGAAA ATTGACCAAG TGATGTGTTT ACATTTACTG AAATGCGCTC
2059

25 TTAAATTTGT TGTAGATTAG GTCTTGCTGG AAGACAGAGA AAAGTTGCCT TTCAGTATTG
2119

ACACTGACTA GAGTGATGAC TGCTTG TAGG TATGTCTGTG CCATTTCTCA GGGAAGTAAG
2179

30 ATGTAAATTG AAGAAGCCTC ACACGTAAAA GAAATGTATT AATGTATGTA GGAGCTGCAG
2239

35 TTCTTG TGGA AGACACTTGC TGAGTGAAGG AAATGAATCT TTGACTGAAG CCGTGCCTGT
2299

AGCCTTG GGG AGGCCCATCC CCCACCTGCC AGCGGTTTCC TGGTGTGGGT CCCTCTGCCC
2359

40 CACCCTCCTT CCCATTGGCT TTCTCTCCTT GGCCTTTCCT GGAAGCCAGT TAGTAAACTT
2419

CCTATTTTCT TGAGTCAAAA AACATGAGCG CTACTCTTGG ATGGGACATT TTTGTCTGTC
2479

5 CTACAATCTA GTAATGTCTA AGTAATGGTT AAGTTTTCTT GTTTCGTCAT CTTTTTGACC
2539

CTCATTCCTT AGAGATGCTA AAATTCCTCG CATAAAGAAG AAGAAATTAA GGAACATAAA
2599

10 TCTTAATACT TGAAGTGTG CCCTTCTGTC CAAGTACTTA ACTATCTGTT CCCTTCCTCT
2659

GTGCCACGCT CCTCTGTTTG TTTGGCTGTC CAGCGATCAG CCATGGCGAC ACTAAAGGAG
2719

15 GAGGAGCCGG GGAAGTCCAG GCTGGAGAGC ACTGCCAGGA CCCACCACTG GAAGCAGGAT
2779

GGAGCTGACT ACGGAAGTGC AACTCAGTG GGCTGTTTCT GCTTATTTCA TCTGTTCTAT
2839

20 GCTTCCTCGT GCCAATTATA GTTTGACAGG GCCTTAAAT TACTTGGCTT TTTCCAAATG
2899

25 CTTCTATTTA TAGAAATCCC AAAGACCTCC ACTTGCTTAA GTATACCTAT CACTTACATT
2959

TTTGTGGTTT TGAGAAAGTA CAGCAGTAGA CTGGGGCGTC ACCTCAGGC CGTTTCTCAT
3019

30 ACTACAGGAT ATTTACTATT ACTCCCAGGA TTCAGCAGAA GATTGCGTTA GCTCTCAAAT
3079

GTGTGTTCCCT GCTTTTCTAA TGGATATTTT AAATTCATT CACAAGCACC TAGTAAGTGC
3139

35 CTGCTGTATC CCTACATTAC ACAGTTCAGC CTTTATCAAG CTTAGTGAGC AGTGAGCACT
3199

40 GAAACATTAT TTTTAAATGT TTAATAAGTT TCTAATATTA AAGTCAGAAT ATTAATACAA
3259

TTAATATTAA TATTAACTAC AGAAAAGACA AACAGTAGAG AACAGCAAAA AAATAAAAAG
3319

5 GATCTCCTTT TTTCCCAGCC CAAATTCTCC TCTCTAAAAG TGTCCACAAG AAGGGGTGTT
3379

TATTCTTCCA ACACATTTCA CTTTTCTGTA AATATACATA AACTTAAAAA GAAAACCTCA
3439

10 TGGAGTCATC TTGCACACAC TTTTCATGCA GTGCTCTTTG TAGCTAAACA GTGAAGATTT
3499

15 ACCTCGTTTCT GCTCAGAGGC CTTGCTGTGG AGCTCCACTG CCATGTACCC AGTAGGGTTT
3559

GACATTTTCAT TAGCCATGCA ACATGGATAT GTATTGGGCA GCAGACTGTG TTTCGTGAAC
3619

20 TGCAGTGATG TATACATCTT ATAGATGCAA AGTATTTTGG GGTATATTAT CCTAAGGGAA
3679

GATAAAGATG ATATTAAGAA CTGCTGTTTC ACGGGGCCCT TACCTGTGAC CCTCTTTGCT
3739

25 GAAGAATATT TAACCCACACA CAGCACTTCA AAGAAGCTGT CTTGGAAGTC TGTCTCAGGA
3799

GCACCCTGTC TTCTTAATTC TCCAAGCGGA TGCTCCATTT CAATTGCTTT GTGACTTCTT
3859

30 CTTCTTTGTT TTTTAAATA TTATGCTGCT TTAACAGTGG AGCTGAATTT TCTGGAAAAT
3919

35 GCTTCTTGGC TGGGGCCACT ACCTCCTTTC CTATCTTTAC ATCTATGTGT ATGTTGACTT
3979

TTTAAAATTC TGAGTGATCC AGGGTATGAC CTAGGGAATG AACTAGCTAT GGAAATAACT
4039

40 CAGGGTTAGG AATCCTAGCA CTTGTCTCAG GACTCTGAAA AGGAACGGCT TCCTCATGCC
4099

TTGTCTTGAT AAAGTGAAT TGGCAAATA GAATTTAGTT TGTACTCAGT GGACAGTGCT
4159

5 GTTGAAGATT TGAGGACTTG TTAAAGAGCA CTGGGTCATA TGGAAAAAAT GTATGTGTCT
4219

CCCCAGGTGC ATTTTCTTGG TTTATGTCTT GTTCTTGAGA TTTTGTATAT TTAGGAAAAC
4279

10 CTCAAGCAGT AATTAATATC TCCTGGAACA CTATAGAGAA CCAAGTGACC GACTCATTTA
4339

CAACTGAAAC CTAGGAAGCC CCTGAGTCCT GAGCGAAAAC AGGAGAGTTA GTCGCCCTAC
4399

15 AGAAAACCCA GCTAGACTAT TGGGTATGAA CTAAAAAGAG ACTGTGCCAT GGTGAGAAAA
4459

ATGTAAAATC CTACAGTGGA ATGAGCAGCC CTTACAGTGT TGTTACCACC AAGGGCAGGT
4519

20 AGGTATTAGT GTTTGAAAAA GCTGGTCTTT GAGCGAGGGC ATAAATACAG CTAGCCCCAG
4579

25 GGGTGGAAACA ACTGTGGGAG TCTTGGGTAC TCGCACCTCT TGGCTTTGTT GATGCTCCGC
4639

CAGGAAGGCC ACTTGTGTGT GCGTGTCTAGT TACTTTTTTA GTAACAATTC AGATCCAGTG
4699

30 TAAACTTCCG TTCATTGCTC TCCAGTCACA TGCCCCCACT TCCCCACAGG TGAAAGTTTT
4759

TCTGAAGTGT TGGGATTGGT TAAGGTCTTT ATTTGTATTA CGTATCTCCC CAAGTCCTCT
4819

35 GTGGCCAGCT GCATCTGTCT GAATGGTGCG TGAAGGCTCT CAGACCTTAC ACACCATTTT
4879

40 GTAAGTTATG TTTTACATGC CCCGTTTTTG AGACTGATCT CGATGCAGGT GGATCTCCTT
4939

GAGATCCTGA TAGCCTGTGA CAGGAATGAA GTAAAGGTCA GTTTTTTTTG TATTGATTTT
4999

5 CACAGCTTTG AGGAACATGC ATAAGAAATG TAGCTGAAGT AGAGGGGACG TGAGAGAAGG
5059

GCCAGGCCGG CAGGCCAACC CTCCTCCAAT GGAAATTCCC GTGTTGCTTC AACTGAGAC
5119

10 AGATGGGACT TAACAGGCAA TGGGGTCCAC TTCCCCCTCT TCAGCATCCC CCGTACC
5176

(2) INFORMATION FOR SEQ ID NO:14:

15

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 451 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

25 Met Asn Leu Glu Gly Gly Gly Arg Gly Gly Glu Phe Gly Met Ser Ala
1 5 10 15

Val Ser Cys Gly Asn Gly Lys Leu Arg Gln Trp Leu Ile Asp Gln Ile
20 25 30

30

Asp Ser Gly Lys Tyr Pro Gly Leu Val Trp Glu Asn Glu Glu Lys Ser
35 40 45

35

Ile Phe Arg Ile Pro Trp Lys His Ala Gly Lys Gln Asp Tyr Asn Arg
50 55 60

Glu Glu Asp Ala Ala Leu Phe Lys Ala Trp Ala Leu Phe Lys Gly Lys
65 70 75 80

40

Phe Arg Glu Gly Ile Asp Lys Pro Asp Pro Pro Thr Trp Lys Thr Arg
85 90 95

	Leu Arg Cys Ala Leu Asn Lys Ser Asn Asp Phe Glu Glu Leu Val Glu	
	100	105 110
5	Arg Ser Gln Leu Asp Ile Ser Asp Pro Tyr Lys Val Tyr Arg Ile Val	
	115	120 125
	Pro Glu Gly Ala Lys Lys Gly Ala Lys Gln Leu Thr Leu Glu Asp Pro	
	130	135 140
10	Gln Met Ser Met Ser His Pro Tyr Thr Met Thr Thr Pro Tyr Pro Ser	
	145	150 155 160
	Leu Pro Ala Gln Gln Val His Asn Tyr Met Met Pro Pro Leu Asp Arg	
		165 170 175
15	Ser Trp Arg Asp Tyr Val Pro Asp Gln Pro His Pro Glu Ile Pro Tyr	
		180 185 190
	Gln Cys Pro Met Thr Phe Gly Pro Arg Gly His His Trp Gln Gly Pro	
20		195 200 205
	Ala Cys Glu Asn Gly Cys Gln Val Thr Gly Thr Phe Tyr Ala Cys Ala	
	210	215 220
25	Pro Pro Glu Ser Gln Ala Pro Gly Val Pro Thr Glu Pro Ser Ile Arg	
	225	230 235 240
	Ser Ala Glu Ala Leu Ala Phe Ser Asp Cys Arg Leu His Ile Cys Leu	
		245 250 255
30	Tyr Tyr Arg Glu Ile Leu Val Lys Glu Leu Thr Thr Ser Ser Pro Glu	
	260	265 270
	Gly Cys Arg Ile Ser His Gly His Thr Tyr Asp Ala Ser Asn Leu Asp	
35		275 280 285
	Gln Val Leu Phe Pro Tyr Pro Glu Asp Asn Gly His Arg Lys Asn Ile	
	290	295 300
40	Glu Asn Leu Leu Ser His Leu Glu Arg Gly Val Val Leu Trp Met Ala	
	305	310 315 320

Pro Asp Gly Leu Tyr Ala Lys Arg Leu Cys Gln Ser Thr Ile Tyr Trp
325 330 335

5 Asp Gly Pro Leu Ala Leu Cys Asn Asp Arg Pro Asn Lys Leu Glu Arg
340 345 350

Asp Gln Thr Cys Lys Leu Phe Asp Thr Gln Gln Phe Leu Ser Glu Leu
355 360 365

10 Gln Ala Phe Ala His His Gly Arg Ser Leu Pro Arg Phe Gln Val Thr
370 375 380

Leu Cys Phe Gly Glu Glu Phe Pro Asp Pro Gln Arg Gln Arg Lys Leu
385 390 395 400

15 Ile Thr Ala His Val Glu Pro Leu Leu Ala Arg Gln Leu Tyr Tyr Phe
405 410 415

Ala Gln Gln Asn Ser Gly His Phe Leu Arg Gly Tyr Asp Leu Pro Glu
20 420 425 430

His Ile Ser Asn Pro Glu Asp Tyr His Arg Ser Ile Arg His Ser Ser
435 440 445

25 Ile Gln Glu
450

(2) INFORMATION FOR SEQ ID NO:15:

30 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 152 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: DNA (genomic)

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

TTTTCTCTAC AGTCACCTCC CTGTTTACCA AAGATAATCA CAATAAGTCC AGTTTACTTA
60

5 CAAAACAAGT TTAGTTATTA GAGGAAACTA AACTTCAGG ATTCAGTCCA GATAATTTTT
120

AAAAACTCTA AAACAATGGA CAGGGCTAGA AT
152

10 (2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 152 base pairs

(B) TYPE: nucleic acid

15 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

25 TGGGCTCGGC CTGGTGGGGC AGCCACAGCG GGACGCAGTA GTGAAAGTCC AGTTTACTTA
60

CAAAACAAGT TTAGTTATTA GAGGAAACTA AACTTCAGG ATTCAGCAGG GCATGAGGAG
120

30

GCAGCTCCTC ACCCTCCCTT TCTCTTTTGT AC
152

(2) INFORMATION FOR SEQ ID NO:17:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 152 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

40

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

5

TGGGCTCGGC CTTGGTGGGG CAGCCACAGC GGGACGCAAG TAGTGAGGGC ACTCAGAACG
60

10

CCACTCAGCC CCGACAGGGC ACTCAGAACG CCACTCAGCC CCGACAGGCA GGGCACGAGG
120

AGGCAGCTCC TCACCCTCCC TTTCTCTTTT GT
152

15

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186

What is claimed is:

1. A method of determining a chromosomal breakpoint in a subject suffering from multiple myeloma which comprises steps of:
 - (a) obtaining a DNA sample from the subject suffering from multiple myeloma;
 - (b) determining whether there is J and C disjunction in the immunoglobulin heavy chain gene in the obtained DNA sample;
 - (c) obtaining a genomic library having clones which contain genomic DNA fragments from the DNA sample which shows positive J and C disjunction;
 - (d) selecting and isolating clones of the obtained library which show positive hybridization with a probe which is capable of specifically hybridizing with the C but not the J region of the immunoglobulin heavy chain gene;
 - (e) preparing fluorescent probes from the genomic DNA fragments of the isolated clones from step (d);
 - (f) hybridizing said fluorescent probes with metaphase chromosomes; and
 - (g) determining the identity of the chromosomes which are capable of hybridizing to said fluorescent probes, wherein the identification of a chromosome other than chromosome 14 would indicate that the chromosomal breakpoint is between chromosome 14 and the identified chromosome, thereby determining a chromosomal breakpoint in a subject suffering from multiple myeloma.
2. The method of claim 1, wherein step (b) is performed by Southern blotting.

3. The method of claim 1, wherein step (b) is performed by polymerase chain reaction with appropriate probes.
4. The method of claim 1, wherein the genomic library is
5 a phage vector library.
5. The method of claim 4, wherein the genomic DNA fragments are generated by cleaving genomic DNA from cells of the subject with an appropriate restriction
10 enzyme.
6. The method of claim 5, wherein the restriction enzyme is *Bam*HI.
7. The method of claim 5, wherein the restriction enzyme
15 is *Sau*3AI.
8. The method of claim 1, wherein the probe of step (d) is a human IgH J region JH probe.
9. The method of claim 1, wherein the probe of step (d) is
20 a human IgH C μ probe.
10. The method of claim 1, wherein the probe of step (d) is
25 a human IgH C γ 2 probe.
11. The method of claim 1, wherein the chromosomal breakpoint identified is a t(6;14)(p25;q32) translocation.
12. The method of claim 1, wherein the chromosomal
30 breakpoint identified is a t(14;15) translocation.

13. A method to identify a gene other than the immunoglobulin gene which is located in chromosome 14, altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma which comprises steps of:

a) selecting a probe having a sequence of a chromosome other than chromosome 14, identified at the chromosomal breakpoint detected in a subject suffering from multiple myeloma, wherein said probe is capable of hybridizing to the unique sequence of the gene other than the immunoglobulin gene altered by a chromosomal breakpoint detected in a subject suffering from multiple myeloma;

b) contacting said probe with mRNA isolated from a cell under conditions permitting formation of a complex between said probe and the mRNA;

c) isolating the complex resulting from step (b);

d) determining the sequence of the mRNA in the isolated complex, thereby determining the identity of the gene.

14. The method of claim 13, wherein step (d) comprises steps of:

i) synthesizing complementary DNA to the mRNA; and

ii) performing sequence analysis of the complementary DNA to determine the sequence of the mRNA.

15. A gene identified by the method of claim 13.

16. The gene of claim 15 designated *MUM-1*.

17. The gene of claim 15 designated *MUM-2*.

18. The method of claim 13, wherein the gene identified comprises a nucleic acid encoding a MUM protein.

19. The method of claim 18, wherein the MUM protein is MUM-1.

20. The method of claim 18, wherein the MUM protein is MUM-2.

21. An isolated nucleic acid molecule encoding a MUM protein.

22. An isolated nucleic acid molecule of claim 21, wherein the nucleic acid molecule is a DNA molecule.

23. The isolated DNA molecule of claim 21, wherein the DNA molecule is a cDNA molecule.

24. The isolated DNA molecule of claim 21, wherein the DNA molecule is a cDNA molecule having the nucleotide sequence shown in Figure 5B (SEQ. ID NO).

25. The isolated DNA molecule of claim 21, wherein the DNA molecule is genomic DNA molecule.

26. The isolated nucleic acid molecule of claim 21, wherein the nucleic acid molecule is an RNA molecule.

5 27. An isolated nucleic acid molecule of claim 21, wherein the nucleic acid molecule encodes a human MUM-1 protein.

10 28. An isolated nucleic acid molecule of claim 21, wherein the nucleic acid molecule encodes a human MUM-2 protein.

15 29. An isolated nucleic molecule of claim 31, wherein the human MUM-1 protein has substantially the same amino acid sequence as shown in Figure 5B (SEQ. ID NO).

30. An isolated nucleic molecule of claim 31, wherein the human MUM-1 protein has the amino acid sequence as shown in Figure 5B (SEQ. ID NO).

20 31. An isolated nucleic acid molecule of claim 21 operatively linked to a promoter of RNA transcription.

25 32. A vector comprising the nucleic acid molecule of claim 21.

33. A vector comprising the nucleic acid molecule of any claim 23.

30 34. A vector comprising the nucleic acid molecule of claim 25.

35. A vector of claim 36, wherein the vector is a plasmid.

36. The plasmid of claim 35, designated pcMUM1-1.6a (ATCC

Accession No.).

37. The plasmid of claim 35, designated pMUM1-2.4B/N (ATCC
Accession No.).

5

38. The plasmid of claim 35, designated pMUM1-7.7B (ATCC
Accession No.).

10

39. The plasmid of claim 35, designated pcMUM2-8 (ATCC
Accession No.).

40. A host cell comprising the vector of claims 32.

15

41. The host cell of claim 40, wherein the cell is selected
from a group consisting of a bacterial cell, a plant
cell, and insect cell and a mammalian cell.

20

42. A nucleic acid probe comprising a nucleic acid molecule
of at least 15 nucleotides capable of specifically
hybridizing with a unique sequence included within the
sequence of a nucleic acid molecule encoding a MUM
protein.

25

43. A nucleic acid probe comprising a nucleic acid molecule
of at least 15 nucleotides which is complementary to a
sequence of the isolated nucleic acid molecule encoding
a MUM protein.

30

44. The nucleic acid probe of either of claims 42 or 43,
wherein the MUM protein is MUM-1.

45. The nucleic acid probe of either of claim 42 or 43,
wherein the MUM protein is MUM-2.

46. A DNA probe of claim 44 or 45.

47. A RNA probe of claim 44 or 45.

5 48. A genomic DNA probe of claim 44 or 45.

49. A nucleic acid probe of claim 44 or 45 labeled with a detectable marker.

10 50. The nucleic acid probe of claim 49, wherein the detectable marker is selected from the group consisting of a radioactive isotope, enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.

15 51. A nucleic acid probe of claim 44, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked to a nucleic acid sequence capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule
20 of human chromosome 14.

52. A nucleic acid probe of claim 45, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked to a nucleic acid sequence capable of
25 specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule of human chromosome 14.

53. A nucleic acid probe comprising a nucleic acid molecule
30 of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule of claim 21 which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

54. The nucleic acid probe of claim 53, wherein the isolated nucleic acid molecule encodes MUM-1.

5 55. The nucleic acid probe of claim 53, wherein the isolated nucleic acid molecule encodes MUM-2.

10 56. The nucleic acid probe of claim 54, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14.

15 57. The nucleic acid probe of claim 55, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14.

20 58. The nucleic acid probe of claim 56, wherein the specific break point comprises a portion of the t(6;14)(p25;q32) translocation.

25 59. The nucleic acid probe of claim 57, wherein the specific break point comprises a portion of a t(14;15) translocation.

30 60. The nucleic acid probe of either of claims 58 or 59 labeled with a detectable marker.

61. The nucleic acid probe of claim 60, wherein the detectable marker is selected from the group consisting of a radioactive isotope, enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.

62. A method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-1

protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-1 protein.

5 63. A method for detecting a predisposition to multiple myeloma associated with the expression of a human MUM-2 protein in a sample from a subject which comprises detecting in a sample from the subject a rearrangement of nucleic acid encoding MUM-2 protein.

10

64. The method of claim 62, wherein the rearrangement of nucleic acid encoding MUM-1 protein is detected by contacting the nucleic acid from the sample with a MUM-1 probe under conditions permitting the MUM-1 probe to hybridize with the nucleic acid encoding MUM-1 protein from the sample, thereby detecting the rearrangement of nucleic acid encoding MUM-1 protein in the sample.

15

65. The method of claim 63, wherein the rearrangement of nucleic acid encoding MUM-2 protein is detected by contacting the nucleic acid from the sample with a MUM-2 probe under conditions permitting the MUM-2 probe to hybridize with the nucleic acid encoding MUM-2 protein from the sample, thereby detecting the rearrangement of nucleic acid encoding MUM-2 protein in the sample.

20

25

66. The method of claim 64, wherein the MUM-1 probe comprises a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-1 protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

30

35 67. The method of claim 65, wherein the MUM-2 probe

comprises a nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-2 protein which is linked to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 15.

68. The method of claim 66, wherein the nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-1 protein is linked at a specific break point to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 14.

69. The method of claim 67, wherein the nucleic acid molecule of at least 15 nucleotides which is complementary to a sequence of the isolated nucleic acid molecule encoding MUM-2 protein is linked at a specific break point to a nucleic acid sequence complementary to a sequence of a nucleic acid molecule of human chromosome 15.

70. The method of claim 68, wherein the specific break point comprises a portion of the t(6;14)(p25;q32) translocation.

71. The method of claim 69, wherein the specific break point comprises a portion of a t(14;15) translocation.

72. The method of claim 62, which comprises:

- a. obtaining DNA from the sample of the subject suffering from multiple myeloma;

- b. performing a restriction digest of the DNA with a panel of restriction enzymes;
- c. separating the resulting DNA fragments by size fractionation;
- d. contacting the resulting DNA fragments with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-1 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 and labeled with a detectable marker;
- e. detecting labeled bands which have hybridized to the nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-1 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 to create a unique band pattern specific to the DNA of subjects suffering from multiple myeloma;
- f. preparing DNA obtained from a sample of a subject for diagnosis by steps (a-e); and
- g. comparing the detected band pattern specific to the DNA obtained from a sample of

subjects suffering from multiple myeloma from step (e) and the DNA obtained from a sample of the subject for diagnosis from step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to multiple myeloma if the patterns are the same.

73. The method of claim 63, which comprises:

- a. obtaining DNA from the sample of the subject suffering from multiple myeloma;
- b. performing a restriction digest of the DNA with a panel of restriction enzymes;
- c. separating the resulting DNA fragments by size fractionation;
- d. contacting the resulting DNA fragments with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-2 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 and labeled with a detectable marker;
- e. detecting labeled bands which have hybridized to the nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-2

protein, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 to create a unique band pattern specific to the DNA of subjects suffering from multiple myeloma;

f. preparing DNA obtained from a sample of a subject for diagnosis by steps (a-e); and

g. comparing the detected band pattern specific to the DNA obtained from a sample of subjects suffering from multiple myeloma from step (e) and the DNA obtained from a sample of the subject for diagnosis from step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to multiple myeloma if the patterns are the same.

74. The method of claim 72 or 73, wherein the size fractionation in step (c) is effected by a polyacrylamide or agarose gel.

75. The method of claim 72 or 73, wherein the detectable marker is radioactive isotope, enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.

76. A method of claim 62 which comprises:

a. obtaining RNA from the sample of the subject suffering from multiple myeloma;

- b. separating the RNA sample by size fractionation;
- c. contacting the resulting RNA species with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-1 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-1 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 14 and labeled with a detectable marker;
- d. detecting labeled bands which have hybridized to the RNA species to create a unique band pattern specific to the RNA of subjects suffering from multiple myeloma;
- f. preparing RNA obtained from a sample of a subject for diagnosis by steps (a-d); and
- g. comparing the detected band pattern specific to the RNA obtained from a sample of subjects suffering from multiple myeloma from step (d) and the RNA obtained from a sample of the subject for diagnosis from step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to multiple myeloma if the patterns are the same.

77. A method of claim 63 which comprises:

- a. obtaining RNA from the sample of the subject

suffering from multiple myeloma;

b. separating the RNA sample by size fractionation;

5

c. contacting the resulting RNA species with a nucleic acid probe capable of specifically hybridizing with a unique sequence included within the sequence of a nucleic acid molecule encoding a human MUM-2 protein, wherein the sequence of a nucleic acid molecule encoding a MUM-2 protein is linked at a specific break point to a specified nucleic acid sequence of human chromosome 15 and labeled with a detectable marker;

10

15

d. detecting labeled bands which have hybridized to the RNA species to create a unique band pattern specific to the RNA of subjects suffering from multiple myeloma;

20

f. preparing RNA obtained from a sample of a subject for diagnosis by steps (a-d); and

25

g. comparing the detected band pattern specific to the RNA obtained from a sample of subjects suffering from multiple myeloma from step (d) and the RNA obtained from a sample of the subject for diagnosis from step (f) to determine whether the patterns are the same or different and to diagnose thereby predisposition to multiple myeloma if the patterns are the same.

30

35 78. The method of claim 76 or 77, wherein the size

fractionation in step (c) is effected by a polyacrylamide or agarose gel.

- 5 79. The method of claim 76 or 77, wherein the detectable marker is radioactive isotope, enzyme, dye, biotin, a fluorescent label or a chemiluminescent label.
- 10 80. The method of either of claim 72 or 76, wherein multiple myeloma associated with the expression of a specific human MUM-1 is diagnosed.
- 15 81. The method of either of claim 73 or 77, wherein multiple myeloma associated with the expression of a specific human MUM-2 is diagnosed.
- 20 82. An antisense oligonucleotide having a sequence capable of specifically hybridizing to an mRNA molecule encoding a human MUM-1 protein so as to prevent overexpression of the mRNA molecule.
- 25 83. An antisense oligonucleotide having a sequence capable of specifically hybridizing to an mRNA molecule encoding a human MUM-2 protein so as to prevent overexpression of the mRNA molecule.
- 30 84. An antisense oligonucleotide having a sequence capable of specifically hybridizing to the cDNA molecule of claim 23.
85. An antisense oligonucleotide having a sequence capable of specifically hybridizing to the genomic DNA molecule of claim 29.
86. An antisense oligonucleotide having a sequence capable

of specifically hybridizing to the RNA molecule of claim 30.

87. An purified MUM protein.

5

88. A purified MUM protein, wherein the MUM protein is MUM-1 protein.

89. A purified human MUM-1 protein of claim 88.

10

90. An antibody directed to a purified MUM-1 protein.

91. An antibody capable of specifically recognizing MUM-1 protein.

15

92. An antibody of claim 91, wherein the MUM-1 protein is a human MUM-1 protein.

93. A purified MUM protein, wherein the MUM protein is MUM-2 protein.

20

94. A purified human MUM-2 protein of claim 93.

95. An antibody directed to a purified MUM-2 protein.

25

96. An antibody capable of specifically recognizing a MUM-2 protein.

97. An antibody of claim 96, wherein the MUM-2 protein is a human MUM-2 protein.

30

98. An monoclonal antibody of any one of claims 90, 91 and 92.

99. An monoclonal antibody of any one of claims 95, 96, and 97.

5 100. A pharmaceutical composition comprising an amount of the oligonucleotide of any one of claims 82, 84, 85 and 81 effective to prevent overexpression of a human MUM-1 protein and a pharmaceutically acceptable carrier capable of passing through a cell membrane.

10 101. A pharmaceutical composition comprising an amount of the oligonucleotide of any one of claims 83, 84, 85 and 81 effective to prevent overexpression of a human MUM-2 protein and a pharmaceutically acceptable carrier capable of passing through a cell membrane.

15

IDENTIFICATION OF GENES ALTERED IN MULTIPLE MYELOMA

Abstract of the Invention

5

10

15


20

25

30

This invention provides a method of determining a chromosomal breakpoint in a subject suffering from multiple myeloma which comprises steps of: (a) obtaining a DNA sample from the subject suffering from multiple myeloma; (b) determining whether there is J and C disjunction in the immunoglobulin heavy chain gene in the obtained DNA sample; (c) obtaining a genomic library having clones which contain genomic DNA fragments from the DNA sample which shows positive J and C disjunction; (d) selecting and isolating clones of the obtained library which show positive hybridization with a probe which is capable of specifically hybridizing with the C but not the J region of the immunoglobulin heavy chain gene; (e) preparing fluorescent probes from the genomic DNA fragments of the isolated clones from step (d); (f) hybridizing said fluorescent probes with metaphase chromosomes; and (g) determining the identity of the chromosomes which are capable of hybridizing to said fluorescent probes, wherein the identification of a chromosome other than chromosome 14 would indicate that the chromosomal breakpoint is between chromosome 14 and the identified chromosome, thereby determining a chromosomal breakpoint in a subject suffering from multiple myeloma. This invention also provides the identified gene altered by a chromosomal breakpoint and various uses thereof.

FIG. 1




14q32 (IgH) germline

FIG. 2A

5kb




der.6
(λSKB-4a)



der.14
(λSKS-3)



6p25 germline



6 TATTAGAGGAACTAAAACTTCAGGATTTCAGTCCAGATAATTTTAAAAACTCTAAACAATGGACAGGGCTAGAAT
|||||
14 TATTAGAGGAACTAAACTTCAGGATTTCAGCAGGGGCATGAGGAGGCAGCTCCTCACCCCTTTCTCTTTGTAC
|||||
14 GGCACCTCAGAACGCCACTCAGCCCCGACAGGCAGGGCACGAGGAGGCAGCTCCTCACCCCTTTCTCTTTGT--
4450 4460 4470 4480 4490 4500 4510

FIG. 3A





FIG. 3B



λ MUM-3


5/21

FIG. 4A



6/21

FIG. 4B



7/21

FIG. 4C




FIG. 5A




FIG. 5B-2

1151 AGAGGGGCGT GGTCCCTCTGG ATGGCCCCCG ACGGGCTCTA TCGGAAAAA
 R G V V L W M A P D G L Y A K R
 1201 CTGTGCCAGA GCACGATCTA CTGGGACGGG CCCCTGGCGC TGTGCAAC
 L C Q S T I Y W D G P L A L C N
 1251 CCGGCCCAAC AAAGTGGAGA GAGACCAGAC CTGCAAGCTC TTTGACAC
 R P N K L E R D Q T C K L F D T
 1301 AGCAGTTCTT GTCAGAGCTG CAAGCGTTTG CTCACCACGG CCGCTCCC
 Q F L S E L Q A F A H H G R S L
 1351 CCAAGATTCC AGGTGACTCT ATGCTTTGGA GAGGAGTTTC CAGACCCT
 P R F Q V T L C F G E E F P D P
 1401 GAGGCAAAGA AAGCTCATCA CAGCTCACGT AGAACCTCTG CTAGCCAG
 R Q R K L I T A H V E P L L A R
 1451 AACTATATTA TTTTGCTCAA CAAAACAGTG GACATTTCCT GAGGGGCT
 L Y Y F A Q Q N S G H F L R S Y
 1501 GATTTACCAG AACACATCAG CAATCCAGAA GATTACCACA GATCTATC
 D L P E H I S N P E D Y H R S I
 1551 CCATTCCCTCT ATTCAAGAAT GAAAAATGTC AAGATGAGTG GTTTTCTT
 H S S I Q E *
 1601 TCCTTTTTTT TTTTTTTTTT TTTGATACGG AGATACGGGG TCTTGCTC
 1651 TCTCCCAGGC TGGAGTGCAG TGACACAATC TCAGCTCACT GTGACCTC
 1701 CCTCCTGGGT TCAAGAGACT CTCCTGCCTC AGCCTCCCTG GTAGCTGG
 1751 TTACAGGTGT GAGCCACTGC ACCCACCCAA GACAAGTGAT TTTCAATTG
 1801 AATATTTGAC TTTAGTGAAA GCGTCCAATT GACTGCCCTC TTAAGTGT
 1851 GAGGAACTCA GAAGTGGAGA TTTGAGTTCA GCGGTTGAGG AGAATTGC
 1901 CGAGACAAGC ATGGAAAATC AGTGACATCT GATTGGCAGA TGAGCTTA
 1951 TCAAAAGGAA GGGTGGCTTT GCATTTTCTT GTGTTCTGTA GACTGCCA
 2001 ATTGATGATC ACTGTGAAAA TTGACCAAGT GATGTGTTTA CATTTACT
 2051 AATGCGCTCT TTAATTTGTT GTAGATTAGG TCTTGCTGGA AGACAGAG
 2101 AACTTGCCTT TCAGTATTGA CACTGACTAG AGTGATGACT GCTTGTAG
 2151 ATGTCTGTGC CATTTCTCAG GGAAGTAAGA TGTAATTGA AGAAGCCT
 2201 CACGTAAAAG AAATGTATTA ATGTATGTAG GAGCTGCAGT TCTTGTGG
 2251 GACACTTGCT GAGTGAAGGA AATGAATCTT TGAAGTGAAG CGTGCCCTG
 2301 GCCTTGGGGA GGCCCATCCC CCACCTGCCA GCGGTTTCCT GGTGTGGG
 2351 CCTCTGCCCC ACCCTCCTTC CCATTGGCTT TCTCTCCTTG GCCTTTCC
 2401 GAAGCCAGTT AGTAAACTTC CTATTTTCTT GAGTCAAAAA ACATGAGC
 2451 TACTCTTGGA TGGGACATTT TTGTCTGTCC TACAATCTAG TAATGTCT
 2501 GTAATGGTTA AGTTTCTTG TTTCTGCATC TTTTGTGACC TCATTCTT
 2551 GAGATGCTAA AATTCTTCGC ATAAAGAAGA AGAAATTAAG GAACATAA
 2601 CTTAATACTT GAACTGTTGC CCTTCTGTCC AAGTACTTAA CTATCTGT
 2651 CCTTCCTCTG TGCCACGCTC CTCTGTTTGT TTGGCTGTCC AGCGATCA
 2701 CATGGCGACA CTAAAGGAGG AGGAGCCGGG GACTCCCAGG CTGGAGAG
 2751 CTGCCAGGAC CCACCACTGG AAGCAGGATG GAGCTGACTA CGGAAGT
 2801 CACTCAGTGG GCTGTTTCTG CTTATTTTCAT CTGTTCTATG CTTCCCTG
 2851 CCAATTATAG TTTGACAGGG CCTTAAAATT ACTTGGCTTT TTCCAAAT
 2901 TTCTATTTAT AGAAATCCCA AAGACCTCCA CTTGCTTAAG TATACCTA
 2951 ACTTACATTT TTGTGGTTTT GAGAAAGTAC AGCAGTAGAC TGGGGCGT

FIG. 5B-3

3051	TCAGCAGAAG	ATTGCGTTAG	CTCTCAAATG	TGTGTTCCCTG	CTTTTCTA
3101	GGATATTTTA	AATTCATTCA	ACAAGCACCT	AGTAAGTGCC	TGCTGTAT
3151	CTACATTACA	CAGTTCAGCC	TTTATCAAGC	TTAGTGAGCA	GTGAGCAC
3201	AAACATTATT	TTTAAATGTT	TAAAAAGTTT	CTAATATTAA	AGTCAGAA
3251	TTAATACAAT	TAATATTAAAT	ATTAACTACA	GAAAAGACAA	ACAGTAGA
3301	ACAGCAAAAA	AATAAAAAGG	ATCTCCTTTT	TTCCCAGCCC	AAATTCTC
3351	CTCTAAAAGT	GTCCACAAGA	AGGGGTGTTT	ATTCTTCCAA	CACATTTT
3401	TTTTCTGTAA	ATATACATAA	ACTTAAAAAG	AAAACCTCAT	GGAGTCAT
3451	TGCACACACT	TTTCATGCAG	TGCTCTTTGT	AGCTAAACAG	TGAAGATT
3501	CCTCGTTCTG	CTCAGAGGCC	TTGCTGTGGA	GCTCCACTGC	CATGTACC
3551	GTAGGGTTTG	ACATTTTCAT	AGCCATGCAA	CATGGATATG	TATTGGGC
3601	CAGACTGTGT	TTCGTGAAC	GCAGTGATGT	ATACATCTTA	TAGATGCA
3651	GTATTTTGGG	GTATATTATC	CTAAGGGAAG	ATAAAGATGA	TATTAAGA
3701	TGCTGTTTCA	CGGGGCCCTT	ACCTGTGACC	CTCTTTGCTG	AAGAATAT
3751	AACCCACAC	AGCACTTCAA	AGAAGCTGTC	TTGGAAGTCT	GTCTCAGG
3801	CACCCGTGCT	TCTTAATTCT	CCAAGCGGAT	GCTCCATTTC	AATTGCTT
3851	TGACTTCTTC	TTCTTTGTTT	TTTTAAATAT	TATGCTGCTT	TAACAGTG
3901	GCTGAATTTT	CTGGAAAATG	CTTCTTGCTT	GGGGCCACTA	CCTCCTTT
3951	TATCTTTACA	TCTATGTGTA	TGTTGACTTT	TTAAAATTCT	GAGTGATC
4001	GGGTATGACC	TAGGGAATGA	ACTAGCTATG	GAAATAACTC	AGGGTTAG
4051	ATCCTAGCAC	TTGTCTCAGG	ACTCTGAAAA	GGAACGGCTT	CCTCATTC
4101	TGTCTTGATA	AAGTGGAATT	GGCAAAC TAG	AATTTAGTTT	GTACTCAG
4151	GACAGTGCTG	TTGAAGATTT	GAGGACTTGT	TAAAGAGCAC	TGGGTCAT
4201	GGAAAAAATG	TATGTGTCTC	CCCAGGTGCA	TTTTCTTGGT	TTATGTCT
4251	TTCTTGAGAT	TTTGTATATT	TAGGAAAACC	TCAAGCAGTA	ATTAATAT
4301	CCTGGAACAC	TATAGAGAAC	CAAGTGACCG	ACTCATTTAC	AACTGAAAC
4351	TAGGAAGCCC	CTGAGTCCTG	AGCGAAAACA	GGAGAGTTAG	TGCCCCTAC
4401	GAAAACCCAG	CTAGACTATT	GGGTATGAAC	TAAAAAGAGA	CTGTGCCAT
4451	GTGAGAAAAA	TGTAAAATCC	TACAGTGGAA	TGAGCAGCCC	TTACAGTG
4501	GTTACCACCA	AGGGCAGGTA	GGTATTAGTG	TTTGAAAAAG	CTGGTCTTT
4551	AGCGAGGGCA	TAAATACAGC	TAGCCCCAGG	GGTGGAACAA	CTGTGGGAC
4601	CTTGGGTACT	CGCACCTCTT	GGCTTTGTTG	ATGCTCCGCC	AGGAAGGCC
4651	CTTGTGTGTG	CGTGT CAGTT	ACTTTTTTAG	TAACAATTCA	GATCCAGTC
4701	AAACTTCCGT	TCATTGCTCT	CCAGTCACAT	GCCCCCACTT	CCCCACAGC
4751	GAAAGTTTTT	CTGAAGTGTT	GGGATTGGTT	AAGGTCTTTA	TTTGTATTA
4801	GTATCTCCCC	AAGTCCTCTG	TGGCCAGCTG	CATCTGTCTG	AATGGTGCC
4851	GAAGGCTCTC	AGACCTTACA	CACCATTTTG	TAAGTTATGT	TTTACATGC
4901	CCGTTTTTTGA	GACTGATCTC	GATGCAGGTG	GATCTCCTTG	AGATCCTGA
4951	AGCCTGTTAC	AGGAATGAAG	TAAAGGTCAG	TTTTTTTTTG	ATTGATTTT
5001	ACAGCTTTGA	GGAACATGCA	TAAGAAATGT	AGCTGAAGTA	GAGGGGACG
5051	GAGAGAAGGG	CCAGGCCGGC	AGGCCAACCC	TCCTCCAATG	GAAATTCCC
5101	TGTTGCTTCA	AACTGAGACA	GATGGGACTT	AACAGGCAAT	GGGGTCCAC
5151	TCCCCCTCTT	CAGCATCCCC	CGTACC		

FIG. 6A

MUM-1 (23-72) KLRQWLIDQI DSGKYPGLMW ENKSI FRI PWKHAGKQDY NREDAALFK
 LSIRF (23-72) KLRQWLIDQI DSGKYPGLMW ENKSV FRI PWKHAGKQDY NREDAALFK
 IRF-1 (7-56) RMRPWLEEQI NSNQIPGLIW INKEMI FQI PWKHAKHGW DINKDACLER
 IRF-2 (7-56) RMRPWLEEQI NSNTIPGLKW INKMKI FQI PWKHAAHGW DVEKDAPLER
 ICSBP (9-60) RLRQWLIEQI DSSMYPGLIW ENKSM FRI PWKHAGKQDY NQLEVDASIEK
 ISGF3Y (11-60) KLRNWWVEQV ESGQFPGVCW DDTAKT M FRI PWKHAGKQDF REDQDAAFK
 IRF-3 (7-55) RILPWLVSQL DLGQLEGVAV VNKSRTR FRI PWKHGTROD. AQQEDFGIEQ

MUM-1 (73-122) AWALEK GKFR EGIDKPD PPT WKTRLRCAIN KSNDFEELVE RSQLDISDPY
 LSIRF (73-122) AWALEK GKFR EGIDKPD PPT WKTRLRCAIN KSNDFEELVE RSQLDISDPY
 IRF-1 (57-106) SWAIHTGRYK AGEKEPD PPT WKANFRCALN SLPTIEELVKI QGRNKGS AV
 IRF-2 (57-106) NWAHTGKHQ PGVDKPD PPT WKANFRCALN SLPTIEELVKI KSIKKGNAF
 ICSBP (59-107) AWALEK GKFK EGDKAEPAT WKTRLRCAIN KSPDFEEVTD RSQLDISEPY
 ISGF3Y (61-109) AWALEK GKYK EGDTGCPAV WKTRLRCAIN KSSEKEVPE RGRMDVAEPY
 IRF-3 (56-104) AWAATGAYV PGRDKPD LPT WKRNFRSALN RKEGLRLAED RSKDPHDPH

MUM-1 (123-130) KVRIVPE
 LSIRF (123-130) KVRIVPE
 IRF-1 (107-114) RVRMIPP
 IRF-2 (107-114) RVRMIPPL
 ICSBP (108-115) KVRIVPE
 ISGF3Y (110-117) KVOQLPP
 IRF-3 (105-112) KIVEFVNS

FIG. 6B

MUM-1 (327-372)	KRLCOSTIYK	DGPLAL....	CNDRENKIER	DOTCKLFDTO	QFLSEDAIA
LSIRF (327-372)	KRLCOSRIYK	DGPLAL....	CSDRENKIER	DOTCKLFDTO	QFLSEDAIA
ICSBP (289-334)	KRLCQGRVFC	SGNAV....	CKGRPNKIER	DEVVQVETS	QFTRFQQY
ISGF3Y (290-335)	QRLCPIPIST	NAPOAP....	PGPGHILIPS	NECVELRFA	YACRDVRYF
IRF-3 (284-333)	QRLGHCHTYK	AVSEELLPNS	GHGPDGEVPK	KEGGVHLG	PLIVDITIT
MUM-1 (373-421)	HHGRSLPRFQ	VTLQFGELFP	DPQRQR.KKI	TAHVPLLLAK	QYYFQQNS
LSIRF (373-421)	HHGRPAPRFQ	VTLQFGELFP	DPQRQR.KKI	TAHVPLLLAK	QYYFQQNT
ICSBP (335-384)	NSQGRIPDGR	VTLQFGELFP	DMAPLRSKHI	LVOIIPOLYR	QIAEEDGKSC
ISGF3Y (336-385)	QGLGPPPKFQ	VTLNMTESH	GSSHTPONFI	TVKMDOAFAN	YILEQTPEQQ
IRF-3 (334-383)	EGSGRSRPRYA	LWFVGEESWP	QDQPWTKRIV	MVKVPTCLR	AVEMRVGG

FIG. 8




FIG. 9A




FIG. 9B




FIG. 10

MUM2 Transcripts detected in MM/PCL Cell Lines




FIG. 11A

Physiological IgH gene rearrangement






FIG. 11B

Chromosomal translocation occurring in switch region



MUM1 cDNA



Sc; SacII, P; *Pst*I, H; *Hind*III, S; *Sac*I, A; *Apa*I

cDNA inserts is cloned into EcoRI / BamHI site of the pBluescript KS+
Bacteria strain used is DH5 α cells. pcMUM1-1.6a contains full length open
reading frame of nt.217-1572.





FIG. 12 A-B

Breakpoint Cloning of the U-266 Cell Line



pMUM2-8 has a 22.0kb insert in BamHI site of pBluescript KS+.

FIG. 13

[illegible]

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below:

<u>Provisional Application No.</u>	<u>Filing Date</u>	<u>Status</u>
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States Application(s), or Section 365(c) of any PCT International Application(s) designating the United States listed below. Insofar as this application discloses and claims subject matter in addition to that disclosed in any such prior Application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56, which became available between the filing date(s) of such prior Application(s) and the national or PCT international filing date of this application:

<u>Application Serial No.</u>	<u>Filing Date</u>	<u>Status</u>
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

And I hereby appoint

John P. White (Reg. No. 28,678); Thomas F. Moran (Reg. No. 16,579); Norman H. Zivin (Reg. No. 25,385); Ivan S. Kavrukov (Reg. No. 25,161); Christopher C. Dunham (Reg. No. 22,031); Thomas G. Carulli (Reg. No. 30,616); Robert D. Katz (Reg. No. 30,141); Peter J. Phillips (Reg. No. 29,691); Richard S. Milner (Reg. No. 33,970); Albert Wai-Kit Chan (Reg. No. 36,479); Kristina L. Konstas (Reg. No. 37,864); Mary Anne P. Tanner (Reg. No. 40,197); Timothy X. Witkowski (Reg. No. 40,232); and Mary Catherine DiNunzio (Reg. No. 37,306)

and each of them, all c/o Cooper & Dunham LLP, 1185 Avenue of the Americas, New York, New York 10036, my attorneys, each with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to receive the patent, to transact all business in the Patent and Trademark Office connected therewith and to file any International Applications which are based thereon under the provisions of the Patent Cooperation Treaty.

Please address all communications, and direct all telephone calls, regarding this application to:

John P. White _____ Reg. No. 28,678
Cooper & Dunham LLP
1185 Avenue of the Americas
New York, New York 10036
Tel. (212) 278-0400

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or
first joint inventor _____ Riccardo Dalla-Favera

Inventor's signature _____ *R. Dalla-Favera*

Citizenship _____ Italy _____ Date of signature _____ 6/6/96

Residence _____ 445 Riverside Drive, #122, New York, New York 10025

Post Office Address _____ same as residence

Full name of joint
inventor (if any) _____

Inventor's signature _____

Citizenship _____ Date of signature _____

Residence _____

Post Office Address _____

Full name of joint
inventor (if any) _____

Inventor's signature _____

Citizenship _____ Date of signature _____

Residence _____

Post Office Address _____

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants : Riccardo Dalla-Favera
U.S. Serial No.: Not Yet Known (Continuation of U.S. Serial
No. 08/654,482, filed May 28, 1996)
Filed : Herewith
For : IDENTIFICATION OF GENES ALTERED IN MULTIPLE
MYELOMA

1185 Avenue of the Americas
New York, New York 10036
June 1, 2000

Assistant Commissioner for Patents
Washington, D.C. 20231

Box: Patent Application

SIR:

ASSOCIATE POWER OF ATTORNEY


Please recognize Jay H. Maioli (Reg. No. 27,213); William E. Pelton
(Reg. No. 25,702); Wendy E. Miller (Reg. No. 35,615); Robert T.
Maldonado (Reg. 38,232); Paul Teng (40,837); Richard F. Jaworski
(Reg. No. 33,515); Elizabeth M. Wieckowski (Reg. No. 42,226);
Pedro C. Fernandez (Reg. No. 41,741); Gary J. Gershik (Reg. No.
39,992); Jane M. Love (Reg. No. 42,812); Spencer H. Schneider (Reg.
No. 45,923); and Raymond A. Diperna (Reg. No. 44,063) as associate
attorney in connection with the above-identified application.

Applicants: Riccardo Dalla-Favera
U.S. Serial No.: Not Yet Known
Filed: Herewith
Page 2

Plase direct correspondence to the attorney of record at the following address:

John P. White
Cooper & Dunham LLP
1185 Avenue of the Americas
New York, New York 10016
(212) 278-0400

Respectfully submitted,

A handwritten signature in black ink, appearing to read "John P. White", written over a horizontal line.

John P. White
Registration No. 28,678
Attorney for Applicant
Cooper & Dunham LLP
1185 Avenue of the Americas
New York, New York 10016
(212) 278-0400